



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/84, 15/82, 5/04, A01H 4/00	A1	(11) International Publication Number: WO 99/07865 (43) International Publication Date: 18 February 1999 (18.02.99)
(21) International Application Number: PCT/US98/16267 (22) International Filing Date: 5 August 1998 (05.08.98) (30) Priority Data: 60/054,836 5 August 1997 (05.08.97) US (71) Applicant: KIMERAGEN, INC. [US/US]; 300 Pheasant Run, Newtown, PA 18940 (US). (72) Inventors: ARNTZEN, Charles, J.; 1005 Highland Road, Ithaca, NY 14850 (US). KIPP, Peter, B.; Apartment 11-3E, 700 Warren Road, Ithaca, NY 14850 (US). KUMAR, Ramesh; 60 Yard Road, Pennington, NJ 08534 (US). MAY, Gregory, D.; 303 The Parkway, Ithaca, NY 14850 (US). (74) Agents: HANSBURG, Daniel; Kimeragen, Inc., 300 Pheasant Run, Newtown, PA 18940 (US) et al.		(81) Designated States: AL, AM, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: THE USE OF MIXED DUPLEX OLIGONUCLEOTIDES TO EFFECT LOCALIZED GENETIC CHANGES IN PLANTS (57) Abstract <p>The invention concerns the use of duplex oligonucleotides about 25 to 30 base pairs to introduce site specific genetic alterations in plant cells. The oligonucleotides can be delivered by mechanical (biolistic) systems or by electroporation of plant protoplasts. Thereafter plants having the genetic alteration can be generated from the altered cells. In specific embodiments the invention concerns alteration in the gene that encodes acid invertase, UDP-glucose pyrophosphorylase, polyphenol oxidase, O-methyl transferase, cinnamyl alcohol dehydrogenase, ACC synthase and ACC oxidase or <i>etr-1</i> or a homolog of <i>etr-1</i>, and plants having isolated point mutations in such genes.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

THE USE OF MIXED DUPLEX OLIGONUCLEOTIDES TO EFFECT LOCALIZED GENETIC CHANGES IN PLANTS

1. FIELD OF THE INVENTION

The field of the present invention relates to methods for the improvement of existing lines of plants and to the development of new lines having desired traits. The previously available methods of obtaining genetically altered plants by recombinant DNA technology enabled the introduction of preconstructed exogenous genes in random, atopic positions, so-called transgenes. In contrast the present invention allows the skilled practitioner to make a specific alteration of a specific pre-existing gene of a plant. The invention utilizes duplex oligonucleotides having a mixture of RNA-like nucleotides and DNA-like nucleotides to effect the alterations, hereinafter "mixed duplex oligonucleotides" or MDON.

2. BACKGROUND TO THE INVENTION

2.1 MDON and Their Use to Effect Specific Genetic Alterations

Mixed duplex oligonucleotides (MDON) and their use to effect genetic changes in eukaryotic cells are described in United States patent No. 5,565,350 to Kmiec (Kmiec I). Kmiec I discloses *inter alia* MDON having two strands, in which a first strand contains two segments of at least 8 RNA-like nucleotides that are separated by a third segment of from 4 to about 50 DNA-like nucleotides, termed an "interposed DNA segment." The nucleotides of the first strand are base paired to DNA-like nucleotides of a second strand. The first and second strands are additionally linked by a segment of single stranded nucleotides so that the first and second strands are parts of a single oligonucleotide chain. Kmiec I further teaches a method for introducing specific genetic alterations into a target gene. According to Kmiec I, the sequences of the RNA segments are selected to be homologous, i.e., identical, to the sequence of a first and a second fragment of the target gene. The sequence of the interposed DNA segment is homologous with the sequence of the target gene between the first and second fragment except for a region of difference, termed the "heterologous region." The heterologous region can effect an insertion or deletion, or can contain one or

more bases that are mismatched with the sequence of target gene so as to effect a substitution. According to Kmiec I, the sequence of the target gene is altered as directed by the heterologous region, such that the target gene becomes homologous with the sequence of the MDON. Kmiec I specifically teaches that ribose and 2'-O-methylribose, i.e., 2'-methoxyribose, containing nucleotides can be used in MDON and that naturally-occurring deoxyribose-containing nucleotides can be used as DNA-like nucleotides.

United States patent application Serial No. 08\664,487, filed June 17, 1996, now U.S. patent No. 5,731,181 (Kmiec II) does specifically disclose the use of MDON to effect genetic changes in plant cells and discloses further examples of analogs and derivatives of RNA-like and DNA-like nucleotides that can be used to effect genetic changes in specific target genes.

Scientific publications disclosing uses of MDON having interposed DNA segments include Yoon, et al., 1996, *Proc. Natl. Acad. Sci.* 93:2071-2076 and Cole-Straus, A. et al., 1996, *SCIENCE* 273 :1386-1389. The scientific publications disclose that rates of mutation as high as about one cell in ten can be obtained using liposomal mediated delivery. However, the scientific publications do not disclose that MDON can be used to make genetic changes in plant cells.

The present specification uses the term MDON, which should be understood to be synonymous with the terms "chimeric mutation vector," "chimeric repair vector" and "chimeraplast" which are used elsewhere.

2.2 Transgenic Plant Cells and the Generation of Plants from Transgenic Plant Cells

Of the techniques taught by Kmiec I and II for delivery of MDON into the target cell, the technique that is most applicable for use with plant cells is the electroporation of protoplasts. The regeneration of fertile plants from protoplast cultures has been reported for certain species of dicotyledonous plants, e.g., *Nicotiana tabacum* (tobacco), United States Patent 5,231,019 and Fromm, M.E., et al., 1988, *Nature* 312, 791, and soybean variety *Glycine max*, WO 92/17598 to Widholm, J.M. However, despite the reports of isolated successes using non-transformed cells, Prioli, L.M., et al., *Bio/Technology* 7, 589, Shillito, R.D., et al., 1989, *Bio/Technology* 7, 581, the regeneration of fertile monocotyledonous plants from transformed protoplast

cultures is not regarded as obtainable with application of routine skill. Frequently, transformed protoplasts of monocotyledonous plants result in non-regenerable tissue or, if the tissue is regenerated the resultant plant is not fertile.

Other techniques to obtain transformed plant cells by introducing kilobase-sized plasmid DNA into plant cells having intact or partially intact cell walls have been developed. United States patent No. 4,945,050, No. 5,100,792 and No. 5,204,253 concern the delivery of plasmids into intact plant cells by adhering the plasmid to a microparticle that is ballistically propelled across the cell wall, hereafter "biolistically transformed" cell. For example U.S. patent No. 5,489,520 describes the regeneration of a fertile maize plant from a biolistically transformed cell. Other techniques for the introduction of plasmid DNA into suspensions of plant cells having intact cell walls include the use of silicon carbide fibers to pierce the cell wall, see U.S. patent No. 5,302,523 to Coffee R., and Dunwell, J.M.

A technique that allows for the electroporation of maize cells having a complex cell wall is reported in U.S. patent No. 5,384,253 to Krzyzek, Laursen and P.C. Anderson. The technique uses a combination of the enzymes endopectin lyase (E.C. 3.2.1.15) and endopolygalacturonase (E.C. 4.2.2.3) to generate transformation competent cells that can be more readily regenerated into fertile plants than true protoplasts. However, the technique is reported to be useful only for F1 cell lines from the cross of line A188 x line B73.

3. SUMMARY OF THE INVENTION

The present invention provides new methods of use of the MDON that are particularly suitable for use in such plant cells.

Thus one aspect of the invention is techniques to adhere MDON to particles which can be projected through the cell wall to release the MDON within the cell in order to cause a mutation in a target gene of the plant cell. The mutations that can be introduced by this technique are mutations that confer a growth advantage to the mutated cells under appropriate conditions and mutations that cause a phenotype that can be detected by visual inspection. Such mutations are termed "selectable mutations."

In a further embodiment the invention encompasses a method of introducing a

mutation other than a selectable mutation into a target gene of a plant cell by a process which includes the steps of introducing a mixture of a first MDON that introduces a selectable mutation in the plant cell and a second MDON that causes the non-selectable mutation.

The invention further encompasses the culture of the cells mutated according to the foregoing embodiments of the invention so as to obtain a plant that produces seeds, henceforth a "fertile plant," and the production of seeds and additional plants from such a fertile plant.

The invention further encompasses fertile plants having novel characteristics which can be produced by the methods of the invention.

4. DETAILED DESCRIPTION OF THE INVENTION

4.1 Recombinagenic Oligonucleobases and Mixed Duplex OligoNucleotides

The invention can be practiced with MDON having the conformations and chemistries described in Kmiec I or in Kmiec II, which are hereby incorporated by reference. The MDON of Kmiec I and/or Kmiec II contain two complementary strands, one of which contains at least one segment of RNA-type nucleotides (an "RNA segment") that are base paired to DNA-type nucleotides of the other strand.

Kmiec II discloses that purine and pyrimidine base-containing non-nucleotides can be substituted for nucleotides. Commonly assigned U.S. patent applications Serial No. 09/078,063, filed May 12, 1998, and Serial No. 09/078,064, filed May 12, 1998, which are each hereby incorporated in their entirety, disclose additional molecules that can be used for the present invention. The term "recombinagenic oligonucleobase" is used herein to denote the molecules that can be used in the present invention. Recombinagenic oligonucleobases include MDON, non-nucleotide containing molecules taught in Kmiec II and the molecules taught in the above noted commonly assigned patent applications.

In a preferred embodiment the RNA-type nucleotides of the MDON are made Rnase resistant by having replacing the 2'-hydroxyl with a fluoro, chloro or bromo functionality or by placing a substituent on the 2'-O. Suitable substituents include the

substituents taught by the Kmiec II, C₁₋₆ alkane. Alternative substituents include the substituents taught by U.S. Patent No. 5,334,711 (Sproat) and the substituents taught by patent publications EP 629 387 and EP 679 657 (collectively, the Martin Applications), which are hereby incorporated by reference. As used herein a 2' - fluoro, chloro or bromo derivative of a ribonucleotide or a ribonucleotide having a 2' - OH substituted with a substituent described in the Martin Applications or Sproat is termed a "2'-Substituted Ribonucleotide." As used herein the term "RNA-type nucleotide" means a 2'-hydroxyl or 2'-Substituted Nucleotide that is linked to other nucleotides of a MDON by an unsubstituted phosphodiester linkage or any of the non-natural linkages taught by Kmiec I or Kmiec II. As used herein the term "deoxyribo-type nucleotide" means a nucleotide having a 2'-H, which can be linked to other nucleotides of a MDON by an unsubstituted phosphodiester linkage or any of the non-natural linkages taught by Kmiec I or Kmiec II.

A particular embodiment of the invention comprises MDON that are linked solely by unsubstituted phosphodiester bonds. Alternatively embodiments comprise linkage by substituted phosphodiesters, phosphodiester derivatives and non-phosphorus-based linkages as taught by Kmiec II. A further particular embodiment comprises MDON wherein each RNA-type nucleotide is a 2'-Substituted Nucleotide. Particular preferred embodiments of 2'-Substituted Ribonucleotides are 2'-fluoro, 2'-methoxy, 2'-propyloxy, 2'-allyloxy, 2'-hydroxylethyloxy, 2'-methoxyethyloxy, 2'-fluoropropyloxy and 2'-trifluoropropyloxy substituted ribonucleotides. In more preferred embodiments of 2'-Substituted Ribonucleotides are 2'-fluoro, 2'-methoxy, 2'-methoxyethyloxy, and 2'-allyloxy substituted nucleotides. In one embodiment the MDON oligomer is linked by unsubstituted phosphodiester bonds.

Although MDON having only a single type of 2'-substituted RNA-type nucleotide are more conveniently synthesized, the invention can be practiced with MDON having two or more types of RNA-type nucleotides. The function of an RNA segment may not be affected by an interruption caused by the introduction of a deoxynucleotide between two RNA-type trinucleotides, accordingly, the term RNA segment encompasses such an "interrupted RNA segment." An uninterrupted RNA segment is termed a contiguous RNA segment. In an alternative embodiment an RNA segment can contain alternating RNase-resistant and unsubstituted 2'-OH nucleotides.

The MDON of the invention preferably have fewer than 100 nucleotides and more preferably fewer than 85 nucleotides, but more than 50 nucleotides. The first and second strands are Watson-Crick base paired. In one embodiment the strands of the MDON are covalently bonded by a linker, such as a single stranded hexa, penta or tetranucleotide so that the first and second strands are segments of a single oligonucleotide chain having a single 3' and a single 5' end. The 3' and 5' ends can be protected by the addition of a "hairpin cap" whereby the 3' and 5' terminal nucleotides are Watson-Crick paired to adjacent nucleotides. A second hairpin cap can, additionally, be placed at the junction between the first and second strands distant from the 3' and 5' ends, so that the Watson-Crick pairing between the first and second strands is stabilized.

The first and second strands contain two regions that are homologous with two fragments of the target gene, i.e., have the same sequence as the target gene. A homologous region contains the nucleotides of an RNA segment and may contain one or more DNA-type nucleotides of connecting DNA segment and may also contain DNA-type nucleotides that are not within the intervening DNA segment. The two regions of homology are separated by, and each is adjacent to, a region having a sequence that differs from the sequence of the target gene, termed a "heterologous region." The heterologous region can contain one, two or three mismatched nucleotides. The mismatched nucleotides can be contiguous or alternatively can be separated by one or two nucleotides that are homologous with the target gene. Alternatively, the heterologous region can also contain an insertion or one, two, three or of five or fewer nucleotides. Alternatively, the sequence of the MDON may differ from the sequence of the target gene only by the deletion of one, two, three, or five or fewer nucleotides from the MDON. The length and position of the heterologous region is, in this case, deemed to be the length of the deletion, even though no nucleotides of the MDON are within the heterologous region. The distance between the fragments of the target gene that are complementary to the two homologous regions is identically the length of the heterologous region when a substitution or substitutions is intended. When the heterologous region contains an insertion, the homologous regions are thereby separated in the MDON farther than their complementary homologous fragments are in the gene, and the converse is applicable

when the heterologous region encodes a deletion.

The RNA segments of the MDON are each a part of a homologous region, i.e., a region that is identical in sequence to a fragment of the target gene, which segments together preferably contain at least 13 RNA-type nucleotides and preferably from 16 to 25 RNA-type nucleotides or yet more preferably 18-22 RNA-type nucleotides or most preferably 20 nucleotides. In one embodiment, RNA segments of the homology regions are separated by and adjacent to, i.e., "connected by" an intervening DNA segment. In one embodiment, each nucleotide of the heterologous region is a nucleotide of the intervening DNA segment. An intervening DNA segment that contains the heterologous region of a MDON is termed a "mutator segment."

Commonly assigned U.S. patent application Serial No. 09/078,063, filed May 12, 1998, and Serial No. 09/078,064, filed May 12, 1998, disclose a type of duplex recombinagenic oligonucleobase in which a strand has a sequence that is identical to that of the target gene and only the sequence of the "complementary" strand contains a heterologous region. This configuration results in one or more mismatched bases or a "heteroduplex" structure. The heterologous region of the heteroduplex recombinagenic oligonucleobases that are useful in the present invention is located in the strand that contains the deoxynucleotides. In one embodiment, the heterologous region is located on the strand that contains the 5' terminal nucleotide.

4.2 The Location and Type of Mutation Introduced by a MDON

Frequently, the design of the MDON for use in plant cells must be modified from the designs taught in Kmiec I and II. In mammalian and yeast cells, the genetic alteration introduced by a MDON that differs from the target gene at one position is the replacement of the nucleotide in the target gene at the mismatched position by a nucleotide complementary to the nucleotide of the MDON at the mismatched position. By contrast, in plant cells there can be an alteration of the nucleotide one base 5' to the mismatched position on the strand that is complementary to the strand that contains the DNA mutator segment. The nucleotide of the target gene is replaced by a nucleotide complementary to the nucleotide of the DNA mutator segment at the mismatched position. Consequently, the mutated target gene differs from the MDON at two positions.

The mutations introduced into the target gene by a MDON are located between the regions of the target gene that are homologous with the ribonucleotide portion of the homology regions of the MDON, henceforth the "RNA segments." The specific mutation that is introduced depends upon the sequence of the heterologous region. An insertion or deletion in the target gene can be introduced by using a heterologous region that contains an insertion or deletion, respectively. A substitution in the target gene can be obtained by using a MDON having a mismatch in the heterologous region of the MDON. In the most frequent embodiments, the mismatch will convert the existing base of the target gene into the base that is complementary to the mismatched base of the MDON. The location of the substitution in the target gene can be either at the position that corresponds to the mismatch or, more frequently, the substitution will be located at the position on the target strand immediately 5' to the position of the mismatch, i.e., complementary to the position of the MDON immediately 3' of the mismatched base of the MDON.

The relative frequency of each location of the mismatch-caused substitution will be characteristic of a given gene and cell type. Thus, those skilled in the art will appreciate that a preliminary study to determine the location of substitutions in the gene of particular interest is generally indicated, when the location of the substitution is critical to the practice of the invention.

4.3 The Delivery of MDON by Microcarriers and Microfibers

The use of metallic microcarriers (microspheres) for introducing large fragments of DNA into plant cells having cellulose cell walls by projectile penetration is well known to those skilled in the relevant art (henceforth biolistic delivery). United States patents No. 4,945,050, No. 5,100,792 and No. 5,204,253 concern general techniques for selecting microcarriers and devices for projecting them.

The conditions that are used to adhere DNA fragments to the microcarriers are not suitable for the use of MDON. The invention provides techniques for adhering sufficient amounts of MDON to the microcarrier so that biolistic delivery can be employed. In a suitable technique, ice cold microcarriers (60 mg/ml), MDON (60 mg/ml) 2.5 M CaCl_2 and 0.1 M spermidine are added in that order; the mixture gently agitated, e.g., by vortexing, for 10 min and allowed to stand at room temperature for

10 min, whereupon the microcarriers are diluted in 5 volumes of ethanol, centrifuged and resuspended in 100% ethanol. Good results can be obtained with a concentration in the adhering solution of 8-10 $\mu\text{g}/\mu\text{l}$ microcarriers, 14-17 $\mu\text{g}/\text{ml}$ MDON, 1.1-1.4 M CaCl_2 and 18-22 mM spermidine. Optimal results were observed under the conditions of 8 $\mu\text{g}/\mu\text{l}$ microcarriers, 16.5 $\mu\text{g}/\text{ml}$ MDON, 1.3 M CaCl_2 and 21 mM spermidine.

MDON can also be introduced into plant cells for the practice of the invention using microfibers to penetrate the cell wall and cell membrane. U.S. Patent No. 5,302,523 to Coffee et al. describes the use of $30 \times 0.5 \mu\text{m}$ and $10 \times 0.3 \mu\text{m}$ silicon carbide fibers to facilitate transformation of suspension maize cultures of Black Mexican Sweet. Any mechanical technique that can be used to introduce DNA for transformation of a plant cell using microfibers can be used to deliver MDON for transmutation.

A suitable technique for microfiber delivery of MDON is as follows. Sterile microfibers (2 μg) are suspended in 150 μl of plant culture medium containing about 10 μg of MDON. A suspension culture is allowed to settle and equal volumes of packed cells and the sterile fiber/MDON suspension are vortexed for 10 minutes and plated. Selective media are applied immediately or with a delay of up to about 120 hours as is appropriate for the particular trait.

The techniques that can be used to deliver MDON to transmute nuclear genes can also be used to cause transmutation of the genes of a plastid of a plant cell. Plastid transformation of higher plants by biolistic delivery of a plasmid followed by an illegitimate recombinatorial insertion of the plasmid is well known to those skilled in the art. Svab, Z., et al., 1990, Proc. Natl. Acad. Sci. **87**, 8526-8530. The initial experiments showed rates of transformation that were between 10-fold and 100-fold less than the rate of nuclear transformation. Subsequent experiments showed that rates of plasmid transformation comparable to the rate of nuclear transformation could be achieved by use of a dominant selectable trait such as a bacterial aminoglycoside 3'-adenosyltransferase gene, which confers spectinomycin resistance. Svab, Z., & Maliga, P., 1993, Proc. Natl. Acad. Sci. **90**, 913-917.

According to the invention MDON for the transmutation of plastid genes can be introduced into plastids by the same techniques as above. When the mutation

desired to be introduced is a selectable mutation the MDON can be used alone. When the desired mutation is non-selectable the relevant MDON can be introduced along with a MDON that introduces a selectable plastid mutation, e.g., a mutation in the *psbA* gene that confers triazine resistance, or in combination with a linear or circular plasmid that confers a selectable trait.

The foregoing techniques can be adapted for use with recombinagenic oligonucleobases other than MDON.

4.4 Protoplast Electroporation

In an alternative embodiment the recombinagenic oligonucleobase can be delivered to the plant cell by electroporation of a protoplast derived from a plant part. The protoplasts are formed by enzymatic treatment of a plant part, particularly a leaf, according to techniques well known to those skilled in the art. See, e.g., Gallois et al., 1996, in *Methods in Molecular Biology* 55, 89-107 (Humana Press, Totowa, NJ). The protoplasts need not be cultured in growth media prior to electroporation.

Suitable conditions for electroporation are 3×10^5 protoplasts in a total volume of 0.3 ml with a concentration of MDON of between 0.6 - 4 $\mu\text{g/mL}$.

4.5 The Introduction of Mutations

The invention can be used to effect genetic changes, herein "transmutate," in plant cells. In an embodiment the plant cells have cell walls, i.e., are other than protoplasts.

The use of MDON to transmutate plant cells can be facilitated by co-introducing a trait that allows for the ready differentiation and separation of cells (hereafter "selection") into which MDON have been introduced from those that have not. In one embodiment of the invention the selection is performed by forming a mixture of MDON and a plasmid that causes the transient expression of a gene that confers a selectable trait, i.e., one that permits survival under certain conditions, e.g., a kanamycin resistance gene. Under these circumstances elimination of cells lacking the selectable trait removes the cells into which MDON were not introduced. The use of a transient expression plasmid to introduce the selectable trait allows for the successive introduction of multiple genetic changes into a plant cell by repeatedly

using a single standardized selection protocol.

In an alternative embodiment transmutation can be used to introduce a selectable trait. A mixture of a first MDON that causes a selectable mutation in a first target gene and a second MDON that causes a non-selectable mutation in a second target gene is prepared. According to the invention, at least about 1% of the cells having the selectable mutation will be found to also contain a mutation in the second target gene that was introduced by the second MDON. More frequently at least about 10% of the cells having the selectable mutation will be found to also contain a mutation in the second target gene.

One use of this embodiment of the invention is the investigation of the function of a gene-of-interest. A mixture is provided of a MDON that causes a selectable mutation and a MDON that causes a mutation that would be expected to "knock-out" the gene-of-interest, e.g., the insertion of a stop codon or a frameshift mutation. Cells in which one or more copies of the gene-of-interest have been knocked out can be recovered from the population having the selectable mutation. Such cells can be regenerated into a plant so that the function of the gene-of-interest can be determined.

A selectable trait can be caused by any mutation that causes a phenotypic change that can produce a selective growth advantage under the appropriate selective conditions or a phenotypic change that can be readily observed, such as change in color of the plant cells growing in a callus. The selectable trait can itself be a desirable traits, e.g., herbicide resistance, or the selectable trait can be used merely to facilitate the isolation of plants having a non-selectable trait that was introduced by transmutation. A desired nonselectable trait can be introduced into a cell by using a mixture of the MDON that causes the desired mutation and the MDON that causes the selectable mutation, followed by culture under the selecting conditions. Selection according to this scheme has the advantage of ensuring that each selected cell not only received the mixture of MDONs, but also that the cell which received the mixture was then susceptible to transmutation by a MDON.

A mutation that causes a lethal phenotypic change under the appropriate conditions, termed a negatively selectable mutation, can also be used in the present invention. Such mutations cause negatively selectable traits. Negatively selectable

traits can be selected by making replica plates of the transmutated cells, selecting one of the replicas and recovering the transmutated cell having the desired property from the non-selected replica.

4.6 Specific Genes That Can Be Transmutated to Create Selectable Traits

In one embodiment of the invention a MDON is used to introduce a mutation into an Acetolactate synthase (ALS) gene, which is also termed the aceto-hydroxy amino acid synthase (AHAS) gene. Sulfonylurea herbicides and imidazoline herbicides are inhibitors of the wild type ALS enzymes. Dominant mutations that render plants resistant to the actions of sulfonylureas and imidazolines have been described. See U.S. Patent Nos. 5,013,659 and 5,378,824 (Bedbrook) and Rajasekaran K., et al., 1996, Mol. Breeding 2, 307-319 (Rajasekaran). Bedbrook at Table 2 describes several mutations (hereafter, a "Bedbrook Mutation") that were found to render yeast ALS enzymes resistant to sulfonylurea herbicides. Bedbrook states that each of the Bedbrook mutations makes a plant resistant to sulfonylurea and imidazoline herbicides when introduced into a plant ALS gene. It is understood that in most plants the gene encoding ALS has been duplicated. A mutation can be introduced into any allele of either ALS gene.

Three of the Bedbrook mutations were, in fact, shown to confer herbicide resistance in a plant, namely the substitutions Pro-Ala¹⁹⁷, Ala-Asp²⁰⁵ and Trp-Leu⁵⁹¹. Rajasekaran reports that mutations Trp-Ser⁵⁹¹ caused resistance to sulfonylurea and imidazoline and that Ser-Asn⁶⁶⁰ caused resistance to imidazoline herbicides. The results of Rajasekaran are reported herein using the sequence numbering of Bedbrook. Those skilled in the art will understand that the ALS genes of different plants are of unequal lengths. For clarity, a numbering system is used in which homologous positions are designated by the same position number in each species. Thus, the designated position of a mutation is determined by the sequence that surrounds it. For example, the mutation Trp-Ser⁵⁹¹ of Rajasekaran is at residue 563 of the cotton ALS gene but is designated as position 591 of Bedbrook because the mutated Trp is surrounded by the sequence that surrounds Trp⁵⁹¹ in Table 2 of Bedbrook. According to the invention any substitution for the naturally occurring amino acid at position 660 or one of the positions listed in Table 2 of Bedbrook, which is hereby incorporated by

reference, can be used to make a selectable mutation in the ALS gene of a plant.

In a further embodiment of the invention the selectable mutation can be a mutation in the chloroplast gene *psbA* that encodes the D1 subunit of photosystem II, see Hirschberg, J., et al., 1984, *Z. Naturforsch.* **39**, 412-420 and Ohad, N., & Hirschberg, J., *The Plant Cell* **4**, 273-282. Hirschberg et al. reports that the mutation Ser-Gly²⁶⁴ results in resistance to triazine herbicides, e.g., 2-Cl-4-ethylamino-6-isopropylamino-s-triazine (Atrazine). Other mutations in the *psbA* gene that cause Atrazine herbicide resistance are described in Erickson J.M., et al., 1989, *Plant Cell* **1**, 361-371, (hereafter an "Erickson mutation"), which is hereby incorporated by reference. The use of the selectable trait caused by an Erickson mutation is preferred when it is desired to introduce a second new trait into a chloroplast.

The scientific literature contains further reports of other mutations that produce selectable traits. Ghislain M., et al., 1995, *The Plant Journal* **8**, 733-743, describes a Asn-Ile¹⁰⁴ mutation in the *Nicotiana sylvestris* dihydrodipicolinate synthase (DHDPs, EC 4.2.1.52) gene that results in resistance to S-(2-aminoethyl)L-cysteine. Mourad, G., & King, J., 1995, *Plant Physiology* **109**, 43-52 describes a mutation in the threonine dehydratase of *Arabidopsis thaliana* that results in resistance to L-O-methylthreonine. Nelson, J.A.E., et al., 1994, *Mol. Cell. Biol.* **14**, 4011-4019 describes the substitution of the C-terminal Leu of the S14/rp59 ribosomal protein by Pro, which causes resistance to the translational inhibitors cryptoluerine and emetine. In further embodiments of the invention, each of the foregoing mutations can be used to create a selectable trait. Each of Ghislain, Mourad and Nelson are hereby incorporated by reference.

4.7 Genes That Can Be Mutated to Create Desirable Non-selectable Traits

Example 1

MALE STERILITY

Certain commercially grown plants are routinely grown from hybrid seed including corn (maize, *Zea mays*), tomatoes and most other vegetables. The production of hybrid seed requires that plants of one purebred line be pollinated only by pollen from another purebred line, i.e., that there be no self pollination. The removal of the pollen-producing organs from the purebred parental plants is a

laborious and expensive process. Therefore, a mutation that induces male-sterility i.e., suppresses pollen production or function, would obviate the need for such process.

Several genes have been identified that are necessary for the maturation or function of pollen but are not essential for other processes of the plant. Chalcone synthase (*chs*) is the key enzyme in the synthesis of flavonoids, which are pigments found in flowers and pollen. Inhibition of *chs* by the introduction of a *chs* antisense expressing gene in the petunia results in male sterility of the plant. Van der Meer, I.M., et al., 1992, *The Plant Cell* **4**, 253-262. There is a family of *chs* genes in most plants. See, e.g., Koes, R.E., et al., 1989, *Plant Mol. Biol.* **12**, 213-226. Likewise disruption of the chalcone synthase gene in maize by insertion of a transposable element results in male sterility. Coe, E.H., *J. Hered.* **72**, 318-320. The structure of maize chalcone synthase and a duplicate gene, *whp*, is given in Franken, P., et al., 1991, *EMBO J.* **10**, 2605-2612. Typically in plants each member of a multigene family is expressed only in a limited range of tissues. Accordingly, the present embodiment of the invention requires that in species having multiple copies of chalcone synthase genes, the particular *chs* gene or genes expressed in the anthers be identified and interrupted by introduction of a frameshift, and one or more in-frame termination codons or by interruption of the promoter.

A second gene that has been identified as essential for the production of pollen is termed *Lat52* in tomato. Muschietti, J., et al., 1994, *The Plant Journal* **6**, 321-338. *LAT52* is a secreted glycoprotein that is related to a trypsin inhibitor. Homologs of *Lat52* have been identified in maize (termed *Zm13*, Hanson D.D., et al., 1989 *Plant Cell* **1**, 173-179; Twell D., et al., 1989, *Mol. Gen. Genet.* **217**, 240-245), rice (termed *Ps1*, Zou J., et al., 1994 *Am. J. Bot.* **81**, 552-561) and olive (termed *Ole e 1*, Villalba, M., et al., 1993, *Eur. J. Biochem.* **276**, 863-869). Accordingly, the present embodiment of the invention provides for a method of obtaining male sterility by the interruption of the *Lat52/Zm13* gene or its homologs by the introduction of a frameshift, one or more in-frame termination codons or by interruption of the promoter.

A third gene that has been identified as essential for the production of pollen is the gene that encodes phenylalanine ammonium lyase (PAL, EC 4.3.1.5). PAL is an essential enzyme in the production of both phenylpropanoids and flavonoids.

Because phenylpropanoids are a precursor to lignins, which can be an essential for the resistance to disease in the preferred embodiment a PAL isozyme that is expressed only in the anther is identified and interrupted to obtain male sterility.

Example 2 ALTERATION OF CARBOHYDRATE METABOLISM OF TUBERS

Once harvested, potato tubers are subject to disease, shrinkage and sprouting during storage. To avoid these losses the storage temperature is reduced to 35-40° F. However, at reduced temperatures, the starch in the tubers undergoes conversion to sugar, termed "cold sweetening", which reduces the commercial and nutritional value of the tuber. Two enzymes are critical for the cold sweetening process: acid invertase and UDP-glucose pyrophosphorylase. Zrenner, R., et al., 1996, *Planta* **198**, 246-252 and Spychalla, J.P., et al., 1994, *J. Plant Physiol.* **144**, 444-453, respectively. The sequence of potato acid invertase is found in EMBL database Accession No. X70368 (SEQ ID NO. 1) and the sequence of the potato UDP Glucose pyrophosphorylase is reported by Katsube, T. et al., 1991, *Biochem.* **30**, 8546-8551. Accordingly, the present embodiment of the invention provides for a method of preventing cold sweetening by the interruption of the acid invertase or the UDP glucose phosphorylase gene by introduction of a frameshift, one or more in-frame termination codons or by interruption of the promoter.

Example 3 REDUCTION IN POST HARVEST BROWNING DUE TO PPO

Polyphenol oxidase (PPO) is the major cause of enzymatic browning in higher plants. PPO catalyzes the conversion of monophenols to o-diphenols and of o-dihydroxyphenols to o-quinones. The quinone products then polymerize and react with amino acid groups in the cellular proteins, which results in discoloration. The problem of PPO induced browning is routinely addressed by the addition of sulfites to the foods, which has been found to be associated with some possible health risk and consumer aversion. PPO normally functions in the defense of the plant to pathogens or insect pests and, hence, is not essential to the viability of the plant. Accordingly, the present embodiment of the invention provides for a method of preventing enzymatic browning by the interruption of the PPO gene by introduction of a frameshift, one or more in-frame termination codons or by interruption of the promoter

in apple, grape, avocado, pear and banana.

The number of PPO genes in the genome of a plant is variable; in tomatoes and potatoes PPO forms a multigene family. Newman, S.M., et al., 1993, *Plant Mol. Biol.* **21**, 1035-1051, Hunt M.D., et al., 1993, *Plant Mol. Biol.* **21**, 59-68; Thygesen, P.W., et al., 1995, *Plant Physiol.* **109**, 525-531. The grape contains only a single PPO gene. Dry, I.B., et al., 1994, *Plant Mol. Biol.*, **26**, 495-502. When the plant species of interest contains multiple copies of PPO genes it is essential that the PPO gene that is normally expressed in the commercial product be interrupted. For example, only one PPO gene is expressed in potatoes of harvestable size, which gene is termed POT32 and its sequence is deposited in GENBANK accession No. U22921 (SEQ ID NO. 2), which sequence is incorporated by reference. The other potato PPO isozymes have been sequenced and the sequences deposited so that one skilled in the art can design a MDON that specifically inactivates POT32.

Example 4 REDUCTION OF LIGNIN IN FORAGE CROPS AND WOOD PULP

Lignin is a complex heterogeneous aromatic polymer, which waterproofs higher plants and strengthens their cell walls. Lignin arises from the random polymerization of free radicals of phenylpropanoid monolignins. Lignins pose a serious problem for the paper industry because their removal from wood pulp involves both monetary and environmental costs. Similarly, the lignin content of forage crops limits their digestibility by ruminants. Indeed, naturally occurring mutations, termed "brown mid-rib" in sorghum, Porter, KS, et al., 1978, *Crop Science* **18**, 205-218, and maize, Lechtenberg, V.L., et al., 1972, *Agron. J.* **64**, 657-660, have been identified as having reduced lignin content and tested as feed for cattle.

The brown mid-rib mutation in maize involves the O-methyl transferase gene. Vignol, F., et al., 1995, *Plant Cell* **7**, 407-416. The O-methyltransferase genes of a number of plant species have been cloned: Burgos, R.C., et al., 1991, *Plant Mol. Biol.* **17**, 1203-1215 (aspen); Gowri, G., et al., 1991, *Plant Physiol.* **97**, 7-14 (alfalfa, *Medicago sativa*) and Jaeck, E., et al., 1992, *Mol. Plant-Microbe Interact.* **4**, 294-300 (tobacco) (SEQ ID No. 3 and SEQ ID No. 4). Thus, one aspect of the present embodiment is the interruption of the O-methyltransferase gene to reproduce a brown mid-rib phenotype in any cultivar of maize or sorghum and in other species of forage

crops and in plants intended for the manufacture of wood pulp.

A second gene that is involved in lignin production is the cinnamyl alcohol dehydrogenase (CAD) gene, which has been cloned in tobacco. Knight, M.E., 1992, *Plant Mol. Biol.* **19**, 793-801 (SEQ ID No. 5 and SEQ ID No. 6). Transgenic tobacco plants making a CAD antisense transcript have reduced levels of CAD and also make a lignin that is more readily extractable, apparently due to an increase in the ratio of syringyl to guaiacyl monomers and to the increased incorporation of aldehyde monomers relative to alcohol residues. Halpin, C., et al., 1994, *The Plant Journal* **6**, 339-350. Accordingly, an embodiment of the invention is the interruption of the CAD gene of forage crops such as alfalfa, maize, sorghum and soybean and of paper pulp trees such as short-leaf pine (*Pinus echinata*) long-leaf pine (*Pinus palustris*) slash pine (*Pinus elliottii*), loblolly pine (*Pinus taeda*), yellow-poplar (*Liriodendron tulipifera*) and cotton wood (*Populus sp.*) by introduction of a frameshift, one or more in-frame termination codons or by interruption of the promoter.

Example 5 THE REDUCTION IN UNSATURATED AND POLYUNSATURATED LIPIDS IN OIL SEEDS

The presence of unsaturated fatty acids, e.g., oleic acid, and polyunsaturated fatty acids, e.g., linoleic and linolenic acids, in vegetable oil from oil seeds such as rape, peanut, sunflower and soybean causes the oils to oxidize, on prolonged storage and at high temperatures. Consequently, vegetable oil is frequently hydrogenated. However, chemical hydrogenation causes transhydrogenation, which produces non-naturally occurring stereo-isomers, which are believed to be a health risk.

Fatty acid synthesis proceeds by the synthesis of the saturated fatty acid on an acyl carrier protein (ACP) followed by the action of desaturases that remove the hydrogen pairs. Consequently, it would be desirable to inhibit the activity of these desaturase enzymes in oil seeds.

The first enzyme in the synthesis of oleic acid is stearoyl-ACP desaturase (EC 1.14.99.6). The stearoyl-ACP desaturases from safflower and castor bean have been cloned and sequenced. Thompson, G.A., et al., 1991, *Proc. Natl. Acad. Sci.* **88**, 2578-2582 (SEQ ID No. 7); Shanklin, J., & Somerville, C., 1991, *Proc. Natl. Acad. Sci.* **88**, 2510-2514 (SEQ ID No. 8); Knutzon, D.S., et al., 1991, *Plant Physiology* **96**, 344-

345. Accordingly, one embodiment of the present invention is the interruption of the stearyl-ACP desaturase gene of oil seed crops such as soybean, safflower, sunflower, soy, maize and rape by introduction of a frameshift, one or more in-frame termination codons or by interruption of the promoter.

A second enzyme that can be interrupted according to the present invention is ω -3 fatty acid desaturase (ω -3 FAD) the enzyme that converts linoleic acid, a diene, to linolenic acid, a triene. There are two ω -3 FAD isozymes in *Arabidopsis thaliana* and, those skilled in the art expect, in most other plants. One isozyme is specific for plastids and is the relevant isozyme for the synthesis of the storage oils of seeds. The other is microsome specific. The cloning of the *Arabidopsis thaliana* plastid ω -3 FAD is reported by Iba., K. et al., 1993, J. Biol. Chem. **268**, 24099-24105 (SEQ ID No. 9). Accordingly an embodiment of the invention is the interruption of the plastid ω -3 FAD gene of oil seed crops such as soybean, safflower, sunflower, soy, maize and rape by introduction of a frameshift, an in-frame termination codon or by interruption of the promoter.

Example 6 INACTIVATION OF S ALLELES TO PERMIT INBRED LINES

Certain plant species have developed a mechanism to prevent self-fertilization. In these species, e.g., wheat and rice, there is a locus, termed S, which has multiple alleles. A plant that expresses an S allele cannot be fertilized by pollen expressing the same S allele. Lee, H-K., et al., 1994, Nature **367**, 560-563; Murfett, J., et al., 1994, Nature **367**, 563. The product of the S locus is an RNase. McClure, B.A., et al., 1989, Nature **342**, 955-957. The product of the S locus is not essential for the plant. Accordingly, an embodiment of the invention is the interruption of genes of the S locus to permit the inbreeding of the plant by introduction of a frameshift, one or more in-frame termination codons or by interruption of the promoter.

Example 7 ETHYLENE INSENSITIVITY

Ethylene is a gaseous plant hormone that is involved in plant growth and development. An unwanted aspect of ethylene's action is the over-ripening of fruit, vegetables and the wilting of flowers that results in rotting and loss. The ethylene

receptor of *Arabidopsis thaliana* has been cloned and is termed ETR-1. Chang, C., et al., 1993, Science **262**, 539-544 (SEQ ID No. 10). A mutant, Cys-Tyr⁶⁵, results in a dominant insensitivity to ethylene. Transgenic tomato plants expressing the *Arabidopsis thaliana* mutant ETR-1 also showed an insensitivity to ethylene, indicating that the Cys-Tyr⁶⁵ mutation would be a dominant suppressor of ethylene action in most plant species. Accordingly one aspect of the present embodiment of the invention is the insertion of the Cys-Tyr⁶⁵ mutation into the ETR-1 gene so as to extend the life span of the mutated fruit vegetable or flower.

In a further aspect of the present embodiment, the preservation of the fruit or flower can be achieved by interrupting one of the genes that encode the enzymes for ethylene synthesis: namely 1-aminocyclopropane-1-carboxylic acid synthase (ACC synthase) and ACC oxidase. For this embodiment of the invention the amount of ethylene synthesis can be eliminated entirely, so that ripening is produced by exogenous ethylene or some amount of ethylene production can be retained so that the fruit ripens spontaneously, but has a prolonged storage life. Accordingly, it is anticipated that the interruption of one allele of either the ACC synthase or the ACC oxidase gene can result in a useful reduction in the level of ethylene synthesis. Alternatively, the invention provides for the interruption of one allele along with the introduction of a mutation that results in a partial loss of activity in the uninterrupted allele.

The sequences of the *Arabidopsis thaliana* ACC synthase and ACC oxidase genes are reported in Abel, S., et al., 1995, J. Biol. Chem. **270**, 19093-19099 (SEQ ID No. 12) and Gomez-Lim, M.A., et al., 1993, Gene **134**, 217-221 (SEQ ID No. 11), respectively, which are incorporated by reference in their entirety.

Example 8

REVERSION OF KANAMYCIN RESISTANCE

Recombinant DNA technology in plants allows for the introduction of genes from one species of plant and bacterial genes into a second species of plant. For example, Kinney, A.J., 1996, Nature Biotech. **14**, 946, describes the introduction of a lauryl ACP-thioesterase gene into the rape seed to obtain a vegetable oil rich in lauric acid. Such transgenic plants are normally constructed using an antibiotic resistance gene, e.g., kanamycin resistance, which is coinserted into the transgenic

plant as a selectable trait. The resultant transgenic plant continues to express the antibiotic resistance gene, which could result in large amounts of the resistance product and the gene entering the food supply and/or the environment, which introduction may represent an environmental or health risk. An embodiment of the invention obviates the risk by providing for the interruption of the kanamycin gene by introduction of a frameshift, one or more in-frame termination codons or by interruption of the promoter.

Example 9 MODIFICATION OF STORAGE PROTEIN AMINO ACID CONTENT

Seeds and tubers contain a family of major storage proteins, e.g., patatins in potato and zeins in maize. The amino acid composition of such storage proteins is often poorly suited to the needs of the human and animals that depend on these crops, e.g., corn is deficient in lysine and methionine and potato is deficient in methionine. Accordingly, one embodiment of the invention is the mutation of a storage protein of a food crop to increase the amount of low abundance amino acids. Patatins are encoded by a multigene family which are characterized in Mignery, G.A., et al., 1988, *Gene* **62**, 27-44, and the structure of zeins is reported by Marks, M.D., et al., 1985, *J. Biol. Chem.* **260**, 16451-459, both of which are hereby incorporated by reference. Alternatively, the anticodon of a methionine or lysine specific tRNA can be mutated to that of a more common amino acid.

Example 10 THE USE OF MDON TO DETERMINE THE FUNCTION OF A GENE

The presently available techniques for the cloning and sequencing of tissue specific cDNAs allow those skilled in the art to obtain readily the sequences of many genes. There is a relative paucity of techniques for determining the function of these genes. In one embodiment of the invention, MDON are designed to introduce frameshift or stop codons into the gene encoding a cDNA of unknown function. This allows for the specific interruption of the gene. Plants having such specific "knock-outs" can be grown and the effects of the knock-out can be observed in order to investigate the function of the unknown gene.

4.8 Fertile Plants of the Invention

The invention encompasses a fertile plant having an isolated selectable point mutation, which isolated selectable mutation is not a rare polymorphism, i.e., would not be found in population of about 10,000 individuals. As used herein a point mutation is mutation that is a substitution of not more than six contiguous nucleotides, preferably not more than three and more preferably one nucleotide or a deletion or insertion from one to five nucleotides and preferably of one or two nucleotides. As used herein an isolated mutation is a mutation which is not closely linked genetically to any other mutation, wherein it is understood that mutations that are greater than 100 Kb and preferably greater than 40 Kb and more preferably greater than 23 Kb are not closely linked.

BIOLISTIC WORKING EXAMPLES

In the following working examples the media and protocols found in Gelvin, S.B., et al., (eds) 1991, PLANT MOLECULAR BIOLOGY MANUAL (Kluwer Acad. Pub.) were followed. Gold particles were coated with MDON according the following protocol. The microprojectiles are first prepared for coating, then immediately coated with the chimeraplast. To prepare the microprojectiles, suspend 60 mg of gold particles in 1 ml of 100% ethanol (see Note 4). Sonicate the suspension for three, 30 s bursts to disperse the particles. Centrifuge at 12,000 xg for 30 s, discard supernatant. Add 1 ml of 100% ethanol, vortex for 15 s, centrifuge at 12,000 xg for 5 min, then discard the supernatant. A 25 μ l suspension of washed gold particles (1.0 μ m diameter, 60 mg/ml) in H₂O are slowly vortexed, to which 40 μ l MDON (50 μ g/ml), 75 μ l of 2.5 M CaCl₂, 75 μ l 0.1M spermidine are sequentially added. All solutions are ice cold. The completed mixture is vortexed for a further 10 min and the particles are allowed to settle at room temperature for a further 10 min. The pellet is washed in 100% EtOH and resuspended in 50 μ l. of absolute ethanol. Biolistic delivery is performed using a Biorad Biolistic gun with the following settings: tank pressure 1100 psi, rupture disks x2 breaking at 900 psi, particle suspension volume 5 μ l.

NT-1 (TOBACCO), A DICOT CELL SUSPENSION: Lawns of NT-1 of approximately 5 cm diameter, containing 5 million cells, were grown for 3 days on standard media at

28°C. Gold particles were coated with ALS-1 or ALS-2 and were shot as above. The cells were cultured a further 2.5 days, suspended and transferred to solid medium supplemented with 15-50 ppb chlorosulfuron (GLEAM™). Resistant colonies emerged after 7-14 days.

The sequences of the MDON used are as follows: (The nucleotides not homologous with the target gene are underlined and bold. Lower case letters denote 2'-O-methyl ribonucleotides.)

ALS-1

```

TGCGCG-guccaguucaCGTTGcauccaacuaT
T                                     T
T                                     T (SEQ ID No. 13)
TCGCGC CAGGTCAAGTGCAACGTAGGATGATT
      3' 5'

```

ALS-2

```

TGCGCG-guccaguucaCGATGcauccaacuaT
T                                     T
T                                     T (SEQ ID No. 14)
TCGCGC CAGGTCAAGTGCTACGTAGGATGATT
      3' 5'

```

ALS-1 and ALS-2 have single base mismatches with the ALS gene at the second nucleotide of the Pro¹⁹⁷ (CCA) codon: ALS-1 is CAA and ALS-2 is CTA. Following PCR amplification and sequencing of the gene of the ALS-1 and ALS-2 transmutated, resistant cell lines, a mutation was in the targeted codon which was found to be Thr (ACA) and Ser (TCA), respectively. The observed mutation was shifted one nucleotide 5' of the location that would have been expected based on the action of MDON in mammalian cells on the coding strand and one nucleotide 3' of the expected location on the non-coding strand. A total of 3 ALS-1 and 5 ALS-2 transmutants having these mutations were identified. No resistant calli were obtained from ALS-1DNA treated cells.

For selection of chlorosulfuron resistant cells, cells were transferred from each bombarded plate to 15 ml containing 5 ml of liquid CSM 2 d after bombardment. The tubes were inverted several times to disperse cell clumps. The cells were then transferred to solidified CSM medium containing 15 ppb chlorosulfuron (Dupont, Wilmington, DE). After approximately 3 - 5 wk, actively growing cells (raised, light

colored colonies) are selected and transferred to solidified CSM containing 50 ppb chlorsulfuron. Three to four weeks later, actively growing cells are selected, then transferred to solidified CSM containing 200 ppb chlorsulfuron. Cells that survive this treatment are then analyzed.

MEDIA

1. NT-1 cell suspension medium (CSM): Murashige and Skoog salts (Gibco BRL, Grand Island, NY), 500 mg/l MES, 1 mg/l thiamine, 100 mg/l myoinositol, 180 mg/l KH_2PO_4 , 2.21 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D), 30g/L sucrose. Adjust pH to 5.7 with 1M KOH or HCl and autoclave. For solidified medium add 8g/l Agar-agar (Sigma, St. Louis, MO) prior to autoclaving.
2. Plating out medium (POM): 80% (v/v) CSM, 0.3M mannitol, 20% (v/v) supernatant from the initial centrifugation of the NT-1 cell suspension prior to protoplast isolation.

TOBACCO LEAF, A DICOT: *Nicotiana tabacum* v. *Samsun* leaf disks were co-transformed by *Agrobacterium tumefaciens* LBA 4404 harboring bin 19-derived plasmids containing a nptII expression cassette containing two genes: a gene for kanamycin resistance and one of two mutants of a gene encoding a Green Fluorescence Protein (GFP, Chui, W., 1996, Current Biol. 6, 325-330). Neither mutant GFP gene produces a GFP product. The mutants contain either a G→T substitution in the sixth codon resulting in a stop codon or a deletion of one nucleotide at the same position, which are termed, respectively, G-stop and G-Δ. After culture on selective MS 104 medium, leaves were recovered and the presence of a GFP gene confirmed by northern blot.

Sequence of first eight codons of GFP:

GFP	ATG GTG AGC AAG GGC GAG GAG CTG	(SEQ ID No. 15)
G-stop	-----T-----	(SEQ ID No. 16)
G-Δ	-----AGG AGC TGT	(SEQ ID No. 17)

The sequences of the MDON used were as follows: (The nucleotides not homologous with G-stop are underlined and bold. Lower case letters denote 2'-O-methyl ribonucleotides.)

GFP - 1

```

TGCGCG-cacucguuccCGCTCcucgacaaguT
T                                     T
T                                     T   (SEQ ID No. 18)
TCGCGC GTGAGCAAGGGCGAGGAGCTGTTCAT
      3' 5'

```

GFP - 2

```

TGCGCG-acucguucccGAGCCucgacaagugT
T                                     T
T                                     T   (SEQ ID NO. 19)
TCGCGC TGAGCAAGGGCTGGGAGCTGTTCACT
      3' 5'

```

Leaf disks of the G-stop and G-Δ transgenic plants were incubated on MS 104 selective media and G-1 or G-1 introduced biologically by two successive deliveries as above. Approximately 10 days after the introduction of the MDON, calli exhibiting GFP-like fluorescence were seen in the G-1 and G-2 treated cultures of both the G-stop and G-Δ leaf disks. Larger and more rapidly growing callusing pieces were subdivided by scalpel to obtain green fluorescent cell-enriched calli. The fluorescent phenotype remained stable for the total period of observation, about 30 days. The presence of green fluorescent cells in the G-1 treated G-stop culture indicates that G-1 does not cause mutations exclusively one base 5' of the mismatched nucleotide.

Green fluorescence was observed using a standard FITC filter set using an IMT-2 Olympus microscope.

ELECTROPORATION WORKING EXAMPLE

CONVERSION OF GFP IN TOBACCO MESOPHYLL PROTOPLASTS

Plant Material

1. Tobacco plant transformant (Delta6) harboring a deletion mutant of GFP.
2. Leaves were harvested from 5 to 6-week-old *in vitro*-grown plantlets

Protoplast Isolation

1. Basically followed the procedure of Gallois, et al., 1996, Electroporation of tobacco leaf protoplasts using plasmid DNA or total genomic DNA. Methods in Molecular Biology, Vol. 55: Plant Cell Electroporation and Electrofusion Protocols Edited by: J. A.

Nickoloff Humana Press Inc., Totowa, NJ. pp.89 - 107.

2. Enzyme solution: 1.2 % cellulase R-10 "Onozuka" (Karlson, Santa Rosa, CA), 0.8 % macerozyme R-10 (Karlson, Santa Rosa, CA), 90 g/l mannitol, 10 mM MES, filter sterilize, store in 10 ml aliquots at -20°C.
3. Leaves were cut from the mid-vein out every 1 - 2 mm. They were then placed abaxial side down in contact with 10 ml of enzyme solution in a 100 x 20 mm petri plate. A total of 1 g of leaves was placed in each plate.
4. The plates were incubated at 25°C in the dark for 16 hr.
5. The digested leaf material was pipetted and sieved through a 100 μ m nylon screen cloth (Small Parts, Inc., Miami Lakes, FL). The filtrate was then transferred to a centrifuge tube, and centrifuged at 1000 rpm for 10 min. All centrifugations for this protocol were done at these conditions.
6. The protoplasts collected in a band at the top. The band of protoplasts was then transferred to a clean centrifuge to which 10 ml of a washing solution (0.4 M sucrose and 80 mM KCl) was added. The protoplasts were gently resuspended, then centrifuged.
7. Repeated step 6 twice.
8. After the last wash, the protoplast density was determined by dispensing a small aliquot onto a hemocytometer. Resuspend the protoplasts to a density of 1×10^6 protoplasts/ml in electroporation buffer (80 mM KCl, 4 mM CaCl_2 , 2mM potassium phosphate, pH 7.2, 8% mannitol, autoclave. The protoplasts were allowed to incubate at 8°C for 2 hr.
9. After 2 hr, 0.3 ml (3×10^5 protoplasts) were transferred to each 0.4 cm cuvette, then placed on ice. GFP-2 (0.6 - 4 μ g/mL) was added to each cuvette except for an unelectroporated control. The protoplasts were electroporated (250V, capacitance 250 μ F, and time constant 10 - 14 ms).
10. The protoplasts were allowed to recover for 10 min on ice, then transferred to petri

plates (100 x 20 mm). After 35 min, 10 ml of POM, see above, was added to each plate. The plates were transferred to the dark at 25°C for 24 hr, then transferred to the light.

11. The protoplast cultures were then maintained according to *Gallois supra*.

Fluorescence Microscopy

1. Under UV light, we observed 8 GFP converted protoplasts out of 3×10^5 protoplasts.

We Claim:

1. A method of making a localized mutation in a target gene in a plant cell comprising the steps of:
 - a. adhering to a particle a recombinagenic oligonucleobase, which contains a first homologous region which has a sequence identical to the sequence of at least 6 base pairs of a first fragment of the target gene and a second homologous region which has a sequence identical to the sequence of at least 6 base pairs of a second fragment of the target gene, and an intervening region which contains at least 1 nucleobase heterologous to the target gene, which intervening region connects the first homologous region and the second homologous region;
 - b. introducing the particle into a cell of a population of plant cells;
 - c. identifying a cell of the population cell having a mutation located between the first and second fragments of the target gene.
2. The method of claim 1, wherein the recombinagenic oligonucleobase is a MDON and each of the homologous regions contains an RNA segment of at least 6 RNA-type nucleotides.
3. The method of claim 2, wherein the intervening region is at least 3 nucleotides in length.
4. The method of claim 2, which further comprises the step of culturing the identified cell so that a plant is generated.
5. The method of claim 2, wherein the first RNA segment contains at least 8 contiguous 2'-Substituted Ribonucleotides.
6. The method of claim 5 wherein the second RNA segment contains at least 8 contiguous 2'-Substituted Ribonucleotides.
7. The method of claim 2, wherein the sequence of the mutated target gene is homologous with the sequence of the MDON.
8. The method of claim 2, wherein the adhering step is performed in a solution

comprising 1.1-1.4 M NaCl and 18-22 μ M spermidine and at least 14 μ g/ml MDON.

9. The method of claim 2, wherein the target gene is a first ALS gene, a second ALS gene, a psbA gene, a threonine dehydratase gene, a dihydrodipicolinate synthase gene, or an S14/rp59 gene
10. The method of claim 9, wherein the plant cell is a maize, wheat, rice or lettuce cell.
11. The method of claim 9, wherein the plant cell is a potato, tomato, canola, soybean or cotton cell.
12. The method of claim 2, wherein the target gene selected from the group consisting of the genes encoding acid invertase, UDP-glucose pyrophosphorylase, polyphenol oxidase, O-methyl transferase, cinnamyl alcohol dehydrogenase, *etr-1* or a homolog thereof, ACC synthase and ACC oxidase.
13. The method of claim 12, where the plant cell is from a maize, wheat, rice or lettuce plant.
14. The method of claim 12, where the plant cell is from a potato, tomato, canola, soybean or cotton plant.
15. The method of claim 2, which further comprises making seeds from the plant or from progeny of the plant.
16. A method of making a localized mutation in a target gene in a plant cell having a cell wall comprising the steps of:
 - a. perforating the cell walls of a population of plant cells;
 - b. introducing a recombinagenic oligonucleobase, which contains a first homologous region which has a sequence identical to the sequence of at least 6 base pairs of a first fragment of the target gene and a second homologous region which has a sequence identical to the sequence of at least 6 base pairs of a second fragment of the target gene, and an intervening region which contains at least 1 nucleobase heterologous to the target gene, which intervening region connects the first homologous region

- and the second homologous region;
- c. identifying a cell of the population having a mutation located between the first and second fragments of the target gene.
17. The method of claim 16, wherein the recombinagenic oligonucleobase is a MDON and each of the homologous regions contains an RNA segment of at least 6 RNA-Type nucleotides.
18. The method of claim 17, which further comprises the step of culturing the identified cell so that a plant is generated.
19. The method of claim 17, wherein the sequence of the target gene between the first and the second fragments differs from the sequence of the intervening region of the MDON at a mismatched nucleotide and the mutation of the target gene is located adjacent to the mismatched nucleotide.
20. The method of claim 17, wherein the sequence of the target gene between the first and the second fragments differs from the sequence of the mutator segment of the MDON at a mismatched nucleotide and the mutation of the target gene is located at the mismatched nucleotide.
21. The method of claim 17, wherein the target gene is a first ALS gene, a second ALS gene, a psbA gene, a threonine dehydratase gene, a dihydrodipicolinate synthase gene, or an S14/rp59 gene
22. The method of claim 21, wherein the plant cell is a maize, wheat, rice or lettuce cell.
23. The method of claim 21, wherein the plant cell is a potato, tomato, canola, soybean or cotton cell.
24. The method of claim 17, wherein the target gene is selected from the group consisting of the genes encoding acid invertase, UDP-glucose pyrophosphorylase, polyphenol oxidase, O-methyl transferase, cinnamyl alcohol dehydrogenase, *etr-1* or a homolog thereof, ACC synthase and ACC oxidase.
25. The method of claim 24, where the target gene is a gene from a maize, wheat, rice or lettuce plant.

26. The method of claim 24, where the target gene is a gene from a potato, tomato, canola, soybean or cotton plant.
27. The method of claim 17, which further comprises making seeds from the plant or from progeny of the plant.
28. A method of making a localized mutation in a target gene of a plastid of a plant cell which comprises the steps of:
 - a. Introducing a recombinagenic oligonucleobase, which contains a first homologous region which has a sequence identical to the sequence of at least 6 base pairs of a first fragment of the target gene and a second homologous region which has a sequence identical to the sequence of at least 6 base pairs of a second fragment of the target gene, and an intervening region which contains at least 1 nucleobase heterologous to the target gene, which intervening region connects the first homologous region and the second homologous region;
 - b. Identifying a cell having a mutation in the region between the first and second fragments of the target gene.
29. The method of claim 28, wherein the recombinagenic oligonucleobase is a MDON and each of the homologous regions contains an RNA segment of at least 6 RNA-Type nucleotides.
30. The method of claim 29, which further comprises culturing the identified cell so that a plant is generated.
31. A method of making a localized, non-selectable mutation in a target gene in a plant cell comprising the steps of:
 - a. introducing into the cells of a population of cells a mixture of a first recombinagenic oligonucleobase and a second recombinagenic oligonucleobase wherein:
 - i. the first recombinagenic oligonucleobase contains a first homologous region which has a sequence identical to the sequence of at least 6 base pairs of a first fragment of a first target gene and a second homologous

- region which has a sequence identical to the sequence of at least 6 base pairs of a second fragment of the first target gene, and an intervening region which contains at least 1 nucleobase heterologous to the target gene, which intervening region connects the first homologous region and the second homologous region, and
- ii. the second recombinagenic oligonucleobase contains a first homologous region which has a sequence identical to the sequence of at least 6 base pairs of a first fragment of a second target gene and a second homologous region which has a sequence identical to the sequence of at least 6 base pairs of a second fragment of the second target gene, and an intervening region which contains at least 1 nucleobase heterologous to the target gene, which intervening region connects the first homologous region and the second homologous region;
 - b. selecting cells from the population having a selectable mutation located between the first and the second fragments of the first target gene from the population; and
 - c. identifying a selected cell having a non-selectable mutation located between the first fragment and the second fragment of the second target cell.
32. The method of claim 31, wherein the each recombinagenic oligonucleobase is a MDON and each of the homologous regions contains an RNA segment of at least 6 RNA-Type nucleotides.
33. The method of claim 32, wherein the first target gene is a first ALS gene, a second ALS gene, a psbA gene, a threonine dehydratase gene, a dihydrodipicolinate synthase gene, or an S14/rp59 gene.
34. The method of claim 33, wherein the plant cell is a maize, wheat, rice or lettuce cell.
35. The method of claim 33, wherein the plant cell is a potato, tomato, canola, soybean or cotton cell.

36. The method of claim 32, wherein the second target gene is selected from the group consisting of the genes encoding acid invertase, UDP-glucose pyrophosphorylase, polyphenol oxidase, O-methyl transferase, cinnamyl alcohol dehydrogenase, *etr-1* or a homolog thereof, ACC synthase and ACC oxidase.
37. The method of claim 36, wherein the plant cell is a maize, wheat, rice or lettuce cell.
38. The method of claim 36, wherein the plant cell is a potato, tomato, canola, soybean or cotton cell.
39. The method of claim 32, which further comprises culturing the identified cell such that a plant is generated.
40. The method of claim 39, which further comprises making seeds from the plant or from progeny of the plant.
41. The method of claim 31, wherein the second recombinogenic oligonucleobase is a heteroduplex recombinogenic oligonucleobase and each of the homologous regions of the second recombinogenic oligonucleobase contains an RNA segment of at least 6 RNA-Type nucleotides.
42. The method of claim 41, wherein the first target gene is a first ALS gene, a second ALS gene, a psbA gene, a threonine dehydratase gene, a dihydrodipicolinate synthase gene, or an S14/rp59 gene.
43. The method of claim 42, wherein the plant cell is a maize, wheat, rice or lettuce cell.
44. The method of claim 42, wherein the plant cell is a potato, tomato, canola, soybean or cotton cell.
45. The method of claim 41, wherein the second target gene is selected from the group consisting of the genes encoding acid invertase, UDP-glucose pyrophosphorylase, polyphenol oxidase, O-methyl transferase, cinnamyl alcohol dehydrogenase, *etr-1* or a homolog thereof, ACC synthase and ACC oxidase..
46. The method of claim 36, 45, wherein the second target gene is from a maize, wheat, rice or lettuce plant.

47. The method of claim 36, 45, wherein the second target gene is from a potato, tomato, canola, soybean or cotton plant.
48. The method of claim 41, which further comprises culturing the identified cell such that a plant is generated.
49. The method of claim 48, which further comprises making seeds from the plant or from progeny of the plant.
50. A method of making a localized mutation in a target gene in a plant cell comprising the steps of:
 - a. digesting a plant part with cellulase such that plant cell protoplasts are formed;
 - b. suspending the protoplasts in a solution comprising a recombinagenic oligonucleobase which contains a first homologous region which has a sequence identical to the sequence of at least 6 base pairs of a first fragment of the target gene and a second homologous region which has a sequence identical to the sequence of at least 6 base pairs of a second fragment of the target gene, and an intervening region which contains at least 1 nucleobase heterologous to the target gene, which intervening region connects the first homologous region and the second homologous region;
 - c. electroporating the suspension such that the recombinagenic oligonucleobase enters a protoplast of the suspension;
 - d. culturing the protoplast; and
 - e. identifying a progeny of the protoplast having a mutation located between the first and second fragments of the target gene.
51. The method of claim 50, which further comprises the step of culturing the identified progeny such that a plant is generated.
52. The method of claim 50, wherein the recombinagenic oligonucleobase is a MDON and each of the homologous regions contains an RNA segment of at least 6 RNA-Type nucleotides.

53. The method of claim 50, wherein the recombinagenic oligonucleobase is an heteroduplex recombinagenic oligonucleobase.
54. A plant or seed having a point mutation in a gene is in its wild type genetic position, which gene is selected from the group consisting of the genes encoding acid invertase, UDP-glucose pyrophosphorylase, polyphenol oxidase, O-methyl transferase, cinnamyl alcohol dehydrogenase, ACC synthase and ACC oxidase or *etr-1* or a homolog of *etr-1*, and the sequence of the genomic DNA within 23 KB of the mutation is the sequence of the wild type DNA, and the point mutation forms a stop codon or is a frameshift mutation.
55. The plant or seed of claim 54, in which the point mutation forms a stop codon.
56. The plant or seed of claim 55, in which the sequence of the genomic DNA within 40 KB of the selectable mutation is the sequence of the wild type DNA.
57. The plant or seed of claim 55, in which the sequence of the genomic DNA within 100 KB of the selectable mutation is the sequence of the wild type DNA.
58. The plant or seed of claim 55, in which the point mutation is a single base pair mutation.
59. The plant or seed of claim 55, which is a maize, wheat, rice or lettuce plant or seed.
60. The plant or seed of claim 55, which is a potato, tomato, canola, soybean or cotton plant or seed.
61. The plant or seed of claim 55, further having a selectable point mutation in a second gene and the sequence of the genomic DNA within 23 KB of the selectable point mutation is the sequence of the wild type DNA.
62. The plant or seed of claim 61, in which the sequence of the genomic DNA within 40 KB of the selectable mutation is the sequence of the wild type DNA.
63. The plant or seed of claim 61, in which the sequence of the genomic DNA within 100 KB of the selectable mutation is the sequence of the wild type DNA.
64. The plant or seed of claim 54, in which the point mutation is a frameshift

mutation.

65. The plant or seed of claim 64, in which the sequence of the genomic DNA within 40 KB of the selectable mutation is the sequence of the wild type DNA.
66. The plant or seed of claim 64, in which the sequence of the genomic DNA within 100 KB of the selectable mutation is the sequence of the wild type DNA.
67. The plant or seed of claim 64, in which the point mutation is a single base pair mutation.
68. The plant or seed of claim 64, which is a maize, wheat, rice or lettuce plant or seed.
69. The plant or seed of claim 64, which is a potato, tomato, canola, soybean or cotton plant or seed.
70. The plant or seed of claim 64, further having a selectable point mutation in a second gene and the sequence of the genomic DNA within 23 KB of the selectable point mutation is the sequence of the wild type DNA.
71. The plant or seed of claim 70, in which the sequence of the genomic DNA within 40 KB of the selectable mutation is the sequence of the wild type DNA.
72. The plant or seed of claim 70, in which the sequence of the genomic DNA within 100 KB of the selectable mutation is the sequence of the wild type DNA.

SEQUENCE LISTING

<110> 1. Arntzen, Charles
 2. Kipp, Peter B.
 3. Kumar, Ramesh
 4. May, Gregory D.

<120> The Use of Mixed Duplex Oligonucleotides
 to Effect Localized Genetic Changes in Plants

<130> 7991-023-999

<150> 60/054,386

<151> 1997-08-05

<160> 19

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 2063

<212> DNA

<213> Solanum tuberosum

<220>

<221> CDS

<222> (3) ... (1907)

<400> 1

agtagcattc	cagttatgac	ccggaaaact	ccgcctccca	ttacacattc	ctcccggatc	60
aaccogattc	cggccaccgg	aagtccctta	aaatcatctc	cggcattttc	ctctcctctt	120
tccttttgct	ttctgtagcc	ttctttccga	tcctcaacaa	ccagtcaccg	gacttgacga	180
gtaactcccg	ttcgccgccc	ccgtcaagag	gtgtttctca	gggagtctcc	gataagact.	240
ttcgagatgt	cgtcaatgct	agtcacattt	cttatgctgt	gtccaatgct	atgcttagct	300
ggcaaaagac	tgcttaccat	tttcaacctc	aaaaaaattg	gatgaacgat	cctaattggc	360
cattgtacca	caagggatgg	tatcatcttt	tttatcaata	caatccagat	tcagctattt	420
ggggaaatat	cacatggggc	catgccgtat	ccaaggactt	gatccactgg	ctctacttgc	480
cttttgccat	ggttcctgat	caatggtagc	atattaacgg	tgtctggact	gggtccgcct	540
ccatcctacc	cgatggtcag	atcatgatgc	tttataccgg	tgtctctgat	gattatgtac	600
aagtgcacaa	tcttgcgtag	cccaccaact	tatctgatcc	tctccttcta	gactgggtca	660
agtacaaaag	caacccggtt	ctggttcctc	caccggcgt	tggtatcaag	gacttttagag	720
acccgaccac	tgcttggtacc	ggaccccaaa	atgggcaatg	gcttttaaca	atcgggtcta	780
agattggtaa	aacgggtatt	gcacttggtt	atgaaacttc	caacttcaca	agctttaagc	840
tattggatga	agtgtgtcat	gcggttccgg	gtacgggtat	gtgggagtgt	gtggactttt	900
accgggtatc	gactgaaaaa	acaaacgggt	tggacacatc	atataacggc	ccgggtgtaa	960
agcatgtgtt	aaaagcaagt	ttagatgaca	ataagcaaga	tcactatgct	attgggacgt	1020
atgacttgac	aaagaacaaa	tggacaccgg	ataaccggga	attggattgt	ggaattgggt	1080
tgaagctgga	ttatgggaaa	tattatgcat	caaagacatt	ttatgacccg	aagaaacaac	1140
gaagagtact	gtggggatgg	attggggaaa	ctgatagtga	atctgctgac	ctgcagaagg	1200
gatgggcatc	tgtacagagt	attccaagga	cagtgtctta	cgacaagaag	acagggacac	1260
atctacttca	gtggccagtt	gaagaaattg	aaagcttaag	agtgggtgat	cctattgtta	1320
agcaagtcaa	tcttcaacca	ggttcaattg	agctactcca	tgttgactca	gctgcagagt	1380
tggatataga	agcctcattt	gaagtggaca	aagtcgcgct	ccagggaata	attgaagcag	1440

```

atcatgtagg tttcagctgc tctactagtg gaggtgctgc tagcagaggc attttggggac 1500
catttgggtg cgttgtaatt gctgatcaaa agctatctga gctaacgcca gtttacttct 1560
acatttctaa aggagctgat ggtcgagctg agactcactt ctgtgctgat caaactagat 1620
cctcagaggc tccgggagtt gctaaacaag tttatggtag ttcagtaccg gtgttagacg 1680
gtgaaaaaca ttcgatgaga ttattggagg accactcaat tgtggagagc tttgcccagg 1740
gaggaagaac agtcataaca tcgcgaattt acccaacaaa ggcagtgaat ggagcagcac 1800
gactcttcgt tttcaacaat gccacagggg ctacgctgac tgcttccgctc aagatttggt 1860
cacttgagtc ggctaataat cgatccttcc ccttgcaaga cttgtaattc atcaagccat 1920
atcttcttca tttttttttt ctttgaagg ttatttcacc gatgtcccat caagaaaggg 1980
aagagagggg gaatatgtag tggtatactc tacttattcg ccatttttagt gatttttcta 2040
ctggactttt gctattcgca aaa 2063

```

```

<210> 2
<211> 1958
<212> DNA
<213> Solanum tuberosum

```

```

<220>
<221> CDS
<222> (22) ... (1815)

```

```

<400> 2
tcttttgcgt tttgagcaat aatggcaagc ttgtgcaata gtagtagtac atctctcaaa 60
actcctttta cttcttcctc cacttcttta tcttccactc ctaagccctc tcaacttttc 120
atccatggaa aacgtaacca aatgttcaaa gtttcatgca aggttaccaa taataacggg 180
gacaaaaacc aaaacgttga aacaaattct gttgatcgaa gaaatgttct tcttggttta 240
gggtggtcttt atggtgttgc taatgctata ccattagctg catccgctgc tccagctcca 300
cctcctgata tctcgtcttg tagtatagcc aggattaacg aaaatcaggg ggtgccgtac 360
agttgttgcg cgcctaagcc tgatgatatg gagaaagttc cgtattacaa gtcccttct 420
atgactaagc tccgtgttcg tcagcctgct catgaagcta atgaggagta tattgccaag 480
tacaatctgg cgattagtcg aatgaaagat cttgataaga cacaaccttt aaaccttatt 540
ggttttaagc aacaagctaa tatacattgt gcttattgta acggtgctta tagaattggg 600
ggcaaagagt tacaagtcca taattcttgg ttttcttcc cgttccatag atggtacttg 660
tacttccacg agagaatcgt gggaaaattc attgatgac caactttcgc tttaccatat 720
tggaattggg accatccaaa aggtatgcgt tttcctgcc tgtatgatcg tgaagggaat 780
tcccttttcg atgtaacacg tgaccaaagt caccgaaatg gagcagtaat cgatcttggg 840
tttttcggca atgaagttga aacaactcaa ctccagttga tgagcaataa tttaacacta 900
atgtaccgtc aaatggtaac taatgctcca tgcctcggg tgttcttttg cgggccttat 960
gatctcgggg ttaacactga actcccggga actatagaaa acatccctca cggctcctgc 1020
cacatctggg ctggtacagt gagaggttca actttgccc atggtgcaat atcaaacggg 1080
gagaatatgg gtcattttta ctacgctggg ttggacccgg ttttcttttg ccatcacagc 1140
aatgtggatc ggatg-ggag cgaatggaaa ggcacaggag ggaaaagaac ggatatcaca 1200
cataaagatt ggttgaactc cgagttcttt ttctatgatg aaaatgaaaa cccttac-gt 1260
gtgaaagtca gagactgttt ggacacgaag aagatgggat acgattacaa accaattgcc 1320
acaccatggc gtaacttcaa gcccttaaca aaggcttcag ctggaaaagt gaatacagct 1380
tcacttccgc cagctagcaa tgtattccca ttggctaaac tcgacaaaagc aatttcgttt 1440
tccatcaata ggccgacttc gtcaaggact caacaagaga aaaatgcaca agaggagatg 1500
ttgacattca gtagcataag atatgataac agagggtaca taagggttca tgtgttttcg 1560
aacgtggaca ataagtgtga tgcgaaatgag cttgacaagg cggagtttgc ggggagttat 1620
acaagtttgc cacatgttca tagagctggg gagactaatc atatcgcgac tgttgatttc 1680
cagctggcga taacggaact gttggaggat attggttttg aagatgaaga tactattgag 1740
gtgactctgg tgccaaagag aggtggtgaa ggtatctcca ttgaagggtc gacgatcagt 1800
cttgacagatt gttaattagt ctctattgaa tctgctgaga ttacactttg atggatgatg 1860
ctctgttttt gttttcttgt tctgtttttt cctctgttga aatcagcttt gttgcttjat 1920
ttcattgaag ttgttattca agaataaatc agttacaa 1958

```

<210> 3
 <211> 1460
 <212> DNA
 <213> Nicotiana tabacum

<220>
 <221> CDS
 <222> (84) ... (1178)

<400> 3

tctgtttctt	caactcacct	taatttgccc	aattgagtea	ttgtaaaatc	tgaaacagaa	60
ccaagagaga	agagaaaaaa	aatatgggtt	caacaagcca	gagccagagt	aagagtctaa	120
ctcacacaga	agacgaagcg	ttcttatttg	ccatgcaatt	ggctagtgtt	tctgtacttc	180
ctatgggtcct	aaaatcagcg	ttagaacttg	accttcttga	actcatggct	aaagctggtc	240
caggtgcagc	cattttctct	tctgaattag	ctgctcagct	ctcaaccag	aaccagaaag	300
caccggttat	tcttgatcgg	atgcttaggc	tacttgctac	ttactctgtt	ctcaattgta	360
ctcttagaac	actgtctgat	ggcagtgttg	agaggcttta	tagtctggct	ccggtttgta	420
agttcttgac	taagaatgct	gatggtgttt	ctgttgcccc	acttttgctt	atgaatcaag	480
ataaagtctt	tatggagagc	tggtagcact	taaaagatgc	agtactagat	ggtggaatcc	540
cattcaacaa	ggcctatgga	atgacagcat	ttgagtacca	tggcacagat	ccaagattca	600
acaaagtgtt	caaccgtgga	atgtctgatc	actccactat	gtcaatgaaa	aagattcttg	660
aggactacaa	aggatttgaa	ggcctaaatt	ccattgttga	tgttggtggg	ggaactggcg	720
ctactgttaa	catgattgtc	tccaaacatc	cctctattaa	gggtattaac	tttgatttac	780
cacatgttat	tggagatgct	ccagcttacc	ctggtgtcga	gcacgttggg	ggcgacatgt	840
ttgccagtgt	gccccaaagca	gatgccattt	tcatagaagt	gatttgtcat	gattggagcg	900
acgagcattg	cctaaaattc	ttgaagaatt	gctatgaagc	actacctgca	aatgggaagg	960
tgataatagc	ggagtgcata	cttccagagg	ccccagatac	atcacttgca	actaagaata	1020
cagtacatgt	tgatattgtg	atgttagcac	ataaccagg	aggcaaagaa	aggactgaga	1080
aggaatttga	ggctttggct	aagggcgctg	gttttactgg	attcgcaagg	cttggtgcgc	1140
ttacaacact	tgggtcatgg	aattcaacaa	ataattaatc	gattcctttg	gaggattaag	1200
caatatactg	ttcattttgc	attttgaat	tctacttttc	acagagtggc	tttactgcga	1260
aataaaagaa	atatatagct	tttaccttga	aaagatcaat	gttcaaaggg	aaaaaaaaaa	1320
aggaagatga	aataattgct	ctcagaaaag	cagtgtgtta	ggaaaaagct	ttttagctgg	1380
attttgaatt	ttattgtatg	tatttctgta	atacacatgt	attgaaggaa	tactagtctt	1440
cgaccaatca	tatttctttg					1460

<210> 4
 <211> 1418
 <212> DNA
 <213> Nicotiana tabacum

<220>
 <221> CDS
 <222> (59) ... (1153)

<400> 4

attccttcaa	cttaccat	taagtcacg	aaaaatctga	aacagaacta	aaagtaaaat	60
gggttcaaca	agcgagagcc	agagtaacag	tctcactcac	acagaagacg	aagctttctt	120
atgtgccatg	caattgtgta	gtgcttctgt	acttcttatg	gtcctaaaat	cagccgtaga	180
acttgacctt	cttgagctaa	tggctaaggc	tggctccagg	gcagctatct	ctccttctga	240
attagctgct	cagctctcaa	ctcagaacct	agaagcacct	gttatgcttg	atcggtatgct	300
taggctactt	gcttcttact	ctgttctcaa	ttgtactctt	agaacactgc	ctgatagcag	360
tggttgagagg	ctttatagtc	tggctcccg	ctgtaagtac	ttgactaaga	atgctgatgg	420
tggttctgtt	gccccacttt	tgcttatgaa	tcaagataaa	gttcttatgg	agagctggta	480
ccacttaaaa	gatgcagtac	tagatggcgg	aatccccattc	aacaaaagcct	atggaatgac	540

4

```

agcatttgag taccatggca cagatccaag attcaacaaa gtgttcaacc gtggaatgtc 600
tgatcactcc actatgtcaa tgaagaagat tcttgaggac tacaaaggat ttgaaggcct 660
aaattccatt gttgatgttg gtggtggaac ggggtgctact gttaacatga ttgtctctaa 720
atatccctct attaagggca ttaactttga tttgccacat gtaattggag atgtccaac 780
ttaccccggt gtcgagcacg ttggtggcga catgtttgct agtgtgccaa aagcagatgc 840
cattttcatg aagtggattt gtcattgattg gtcgagatgag cattgcctaa aattcttgaa 900
gaattgctat gaagcactac ctgcaaattg gaaggtgata attgcagagt gcataacttc 960
agaggcccca gatacatcac ttgcaactaa gaatacagta catgttgata ttgttatgtt 1020
agcacataac ccaggaggca aagaaaggac tgagaaggaa tttgaggctt tggctaaggg 1080
cgctggtttt actggattcg caaggcttgt tgcgcttaca acacttgggt catggaatc 1140
aacaagtaat taatcgattc cttaatttga aggattaagc aatatactgt tcgttttgca 1200
tttgaaatt ctacttttct cagagtggct tgactgtgaa ataaagaaa tatagctttt 1260
aactgaaaa gattgatgtt caaaagaaaa aaaggaagat gaaataattg ctctcagaaa 1320
agcaatgtgt taggaaaaag ctttttttagc tggattttga attttactgt atgtatttct 1380
gttatacaca tgtattgaag gaatactagt ttcgcacc 1418

```

<210> 5

<211> 1419

<212> DNA

<213> Nicotiana tabacum

<220>

<221> CDS

<222> (92) ... (1165)

<400> 5

```

atttctttct ctttcccttg aactgtgttt tcattttttc tgctctgaaa caatagtgtt 60
ttccttgtag attttaagtt aaaagaaaaac catgggtagc ttggatgttg aaaaatcagc 120
tattgggttg gctgctagag acccttcttg tctactttca ccttatacct atactctcag 180
aaacacagga cctgaagatg tgcaagtcaa agttttgtat tgtggacttt gccacagtga 240
tcttcaccaa gttaaaaatg atcttggcat gtccaactac cctctggttc ctggacatga 300
agtgggtgga aaagtagtgg aggtaggagc agatgtgtca aaattcaaaag tgggggacac 360
agttggagtt ggattactcg ttggaagttg taggaactgt ggcccttgca agagagaaat 420
agagcaatat tgcaacaaga agatttggaa ttgcaatgat gtctacactg atggcaaacc 480
caccacaagg gtgttttgcta attctatggg tgttgatcaa aactttgttg tgaaaattcc 540
agaggggtatg gcaccagaac aagcagcacc tctattatgt gctggcataa cagtatacag 600
tccattcaac catttttggt ttaatcagag tggatttaga ggaggaattt tgggattagg 660
aggagtggga catatgggag tgaaaatagc aaaggcaatg ggacatcatg ttactgtcat 720
tagttcttca aataagaaga gacaagaggc attggaacat cttggtgcag atgattatct 780
tgtagttca gacactgata aaatgcaaga agctgctgat tcacttgact atattattga 840
tactgtccct gttggccatc ctcttgaaact ttatctttct ttgcttaaaa ttgatggcaa 900
acttatcttg atcggagtta tcaacacccc cttgcaattt atctctccca tgggttatgt 960
cgggagaaaag agcatcactg gaagctttat tggtagcatg aaggaaacag aggaaatgct 1020
agacttctgc aaagagaaaag gtgtgacttc acagattgag atagtgaaaa tggattatat 1080
caacactgca atggagaggt tggagaaaaa tgatgtgagc tacagatttg ttgttgatgt 1140
tgctggaagc aagcttgacc agtaattgca caagaaaaac aacatggaat ggttcactat 1200
tatacaacaa ggctatgaga aaaatagtac tctcaactt tgatgtcatc ttgtttacct 1260
ttgttttatt tccacctgt attatcatat ttgggtggcgt agagtgcagt ttatgtatat 1320
tttctttctt caaaacaatc ttaaatgaat ttggatgttg gtgacgattt tgaaatatac 1380
caaccatgca aacttacttt ggtagaaaaa aaaaaaaaaa 1419

```

<210> 6

<211> 1398

<212> DNA

<213> Nicotiana tabacum

<220>
 <221> CDS
 <222> (88) ... (1161)

<400> 6

attcctcctt	cccttgaact	gtgttttctg	tttttctgct	ctaaaacaat	cgtgtgttcc	60
ttctagattt	taagttttaa	gaacatcatg	ggtggcttgg	aagttgagaa	aacaactatt	120
ggttgggctg	ctagagaccc	ttctgggtga	ctttcacctt	atacctatac	tctcagaaac	180
acaggacctg	aagatgtgga	agtcaaagtt	ttgtattgtg	ggctctgtca	cactgatctt	240
caccaagtta	aaaatgatct	tggcatgtcc	aactaccctc	tggttcctgg	acatgaagtg	300
gtgggagaag	tgggtggaggt	aggaccagat	gtgtcaaaat	tcaaagttgg	ggacacagtt	360
ggagtggat	tactcgttgg	aagttgcagg	aactgtggcc	cttgcaagag	agatatagag	420
caatattgca	acaagaagat	ttggaactgc	aatgatgtct	acactgatgg	caaaccacc	480
caaggtgggt	ttgctaaatc	catggttgtt	gatcaaaagt	ttgtggtgaa	aattccagag	540
ggtatggcac	cagaacaagc	agcacctcta	ttatgtgctg	gtataacagt	atacagtcca	600
ttgaaccatt	ttggtttcaa	acagagtggg	ttaagaggag	gaattttggg	attaggaggga	660
gtgggacaca	tgggagtga	aatagcaaag	gcaatgggac	atcatgttac	tgtcattagt	720
tcttcaaata	agaagagaca	agaggcattg	gaacatcttg	gtgcagatga	ttatcttgtc	780
agttcagaca	ctgataaaat	gcaagaggct	tctgattcac	ttgactatat	tattgatact	840
gtccctgttg	gccatcctct	tgaaccttat	ctttctttgc	ttaaaattga	tggcaaacct	900
atcttgatgg	gagttatcaa	cacccctctg	caatcttatct	cccccatggt	tatgctcggg	960
agaaagagca	tcacaggaag	ctttattgggt	agcatgaagg	aaacagaggga	aatgctagat	1020
ttctgcaaag	agaaagggtg	gacttcacag	attgagatag	tgaaaatgga	ttatatcaac	1080
actgcaatgg	agagggttga	gaaaaatgat	gtgaggtaca	gatttgtggg	tgatgttatt	1140
ggaagcaagc	ttgaccagta	attatattac	acaagaaaaa	caacatggaa	tggttcacta	1200
ttatacaagg	ctgtgagaat	actaaacttt	gatgtcgtct	tttgtatcct	tttgttttat	1260
ttgccacctg	tattttctta	tttgggtgatc	gagagtgcag	tttatgtatt	attttctttc	1320
ttcaaaacaa	tttaatgtat	gaatttggat	gttgggtgaa	aaaaaaaaaa	aaaaaaaaaa	1380
aaaaaaaaaa	aaaaaaaaaa					1398

<210> 7
 <211> 1533
 <212> DNA
 <213> *Carthamus tinctorius*

<220>
 <221> CDS
 <222> (106) ... (1296)

<400> 7

gctcacttgt	gtgggtggagg	agaaaaacag	aactcacaaa	aagctttgctg	actgccaaaga	60
acaacaacaa	caacaagatc	aagaagaaga	agaagaagat	caaaaatggc	tcttcgaatc	120
actccagtga	ccttgcaatc	ggagagatac	cgttcgtttt	cgtttcctaa	gaaggctaata	180
ctcagatctc	ccaaattcgc	catggcctcc	accctcggat	catccacacc	gaagggtgac	240
aatgccaaaga	agccttttca	acctccacga	gagggttcatg	ttcagggtgac	gcactccatg	300
ccaccacaga	agatagagat	tttcaaattc	atcgagggtt	gggctgagca	gaacatattg	360
gttcacctaa	agccagtggg	gaaatgttgg	caagcacagg	atttcttgcc	ggaccctgca	420
tctgaaggat	ttgatgaaca	agtcaaggaa	ctaagggcaa	gagcaaagga	gattcctgat	480
gattactttg	ttgttttggg	tggagatatg	attacagagg	aagccctacc	tacttaccaa	540
acaatgctta	ataccctaga	tgggtgtacgt	gatgagactg	gggctagcct	tacgccttgg	600
gctgtctgga	ctagggtctg	gacagctgaa	gagaacaggc	atggcgatct	tctccacacc	660
tatctctacc	tttctgggctg	ggtagacatg	aggcagatac	agaagacaat	tcagtatctc	720
attgggtcag	gaatggatcc	tcgtaccgaa	aacagcccct	accttggggt	catctacaca	780
tcgtttcaag	agcgtgccac	atttgtttct	cacggaaaca	ccgccaggca	tgcaaaagga	840
catggggacg	tgaactggc	gcaaatttgt	ggtacaatcg	cgtctgacga	aaagcgtcac	900

```

gagaccgctt atacaaagat agtcgaaaag ctattcgaga tcgatacctga tggcaccgtt      960
cttgcttttg ccgacatgat gaggaaaaag atctcgatgc ccgcacactt gatgtacgat      1020
gggcgtgatg acaacctctt cgaacatttc tcggcgggtg cccaaagact cggcgtctac      1080
accgccaaag actacgccga catactggaa tttctggctg ggcgggtggaa agtggcggat      1140
ttgaccggcc tatctggtga agggcgtaaa gcgcaagatt atgtttgcg gttgccacca      1200
agaatcagaa ggctggagga gagagctcaa gggcgagcaa aggaaggacc tgttgttcca      1260
ttcagctgga ttttcgatag acaggtgaag ctgtgaagaa aaaaaaacg agcagtgagt      1320
tcggtttctg ttggcttatt gggtagaggt taaaacctat tttagatgtc tgtttcgtgt      1380
aatgtggttt tttttcttct aatcttgaat ctggtattgt gtcggtgagt tcgctgtgtg      1440
gtaaaacttg gtggctgtgg acatattata gaactcgta tgccaatttt gatgacggtg      1500
gttatcgctt cccctggtgt ttttttattg ttt                                     1533

```

<210> 8

<211> 1643

<212> DNA

<213> Ricinus communis

<220>

<221> CDS

<222> (1) ... (1239)

<400> 8

```

ttccggcaaa taacaaaaaa ccaaaagaaa aaggtaagaa aaaaaacaat ggctctcaag      60
ctcaatcctt tccttttctc aaccctaaaag ttaccttctt tcgctcttcc accaatggcc      120
agtaccagat ctccctaagt ctacatggcc tctacctca agtctggttc taaggaaagt      180
gagaatctca agaagccttt catgcctcct cgggagggtac atgttcaggt taccattctt      240
atgccacccc aaaagattga gatctttaaa tccctagaca attgggctga ggagaacatt      300
ctggttcctc tgaagccagt tgagaaatgt tggcaaccgc aggatttttt gccagatccc      360
gcctctgatg gatttgatga gcaagtcagg gaactcaggg agagagcaaa ggagattcct      420
gatgattatt ttgttggttt gggtggagac atgataacgg aagaagccct tcccacttat      480
caaacaatgc tgaatacctt ggatggagtt cgggatgaaa cagggtgcaag tcctacttct      540
tgggcaattt ggacaagggc atggactgctg gaagagaata gacatggtga cctcctcaat      600
aagtatctct acctatctgg acgagtggac atgaggcaaa ttgagaagac aattcaataa      660
ttgattggtt caggaatgga tccacggaca gaaaacagtc cataccttgg gttcatctat      720
acatcattcc aggaaagggc aaccttcatt tctcatggga aactgcccg acaagccaaa      780
gagcatggag acataaagt ggctcaaata tgtggtacaa ttgctgcaga tgagaagcgc      840
catgagacag cctacacaaa gatagtggaa aaactctttg agattgatcc tgatggaact      900
gttttggtct ttgctgatat gatgagaaag aaaatcttct tgctgcaca cttgatgtat      960
gatggccgag atgataatct ttttgaccac ttttcagctg ttgcgcagcg tcttgagtc      1020
tacacagcaa aggattatgc agatatattg gagttcttgg tgggcagatg gaaggtggat      1080
aaactaacgg gcctttcagc tgagggacaa aaggctcagg actatgtttg tcggttacct      1140
ccaagaatta gaaggtcgga agagagagct caaggaaggg caaaggaagc acccaccatg      1200
cctttcagct ggattttcga taggcaagtg aagctgtagg tggctaaagt gcaggacgaa      1260
accgaaatgg ttagtttcac tctttttcat gccatccct gcagaatcag aagtagaggt      1320
agaattttgt agttgctttt ttattacaag tccagtttag tttaaggctc gtggaaggga      1380
gttagttgag gagtgaatct agtaagttgt tgatactggt gtgttcttgt gttgtcatga      1440
gtctgcttga tagtgagttt cttttgttct cttttgttgt gttcttttat ctggtctctc      1500
tctctctctc tctctctttt tctcttacc caagtgtctc aagtataata agcaaacgat      1560
ccatgtggca attttgatga tggatgatcag tctcacaact tgatcttttg tcttctattg      1620
gaaacacagc ctgcttggtt gaa                                     1643

```

<210> 9

<211> 2569

<212> DNA

<213> Arabidopsis thaliana

```

<220>
<221> exon
<222> (236) ... (729)
<223> Exon 1

<221> exon
<222> (1030) ... (1119)
<223> Exon 2

<221> exon
<222> (1201) ... (1267)
<223> Exon 3

<221> exon
<222> (1358) ... (1450)
<223> Exon 4

<221> exon
<222> (1530) ... (1715)
<223> Exon 5

<221> exon
<222> (1809) ... (1889)
<223> Exon 6

<221> exon
<222> (1993) ... (2130)
<223> Exon 7

<221> exon
<222> (2212) ... (2403)
<223> Exon 8

<400> 9
cacaccatca ctaataaatt tccttctcct ttcaagttgt agctaactta tataagacat      60
aagcgtgcga accagagaca gagatagaaa ttgagagacg ataagcaaag tagaaaacac      120
aagtttctct cacacacatt atctctttct ctattaccac cactcattca taacagaaac      180
ccacaaaaaa ataaaaagag agacttttca ctctggggag agagctcaag ttctaattggc      240
gaacttggtc ttatcagaat gtggtatacg acctctcccc agaatctaca caacaccag      300
atccaatttc ctctccaaca acaacaaatt cagaccatca ctttcttctt cttcttataa      360
aacatcatca tctcctctgt cttttggtct gaattcacga gatgggttca cgaggaattg      420
ggcgttgaat gtgagcacac cattaacgac accaatattt gaggagtctc cattggagga      480
agataataaa cagagattcg atccagggtgc gcctcctccg ttcaatttag ctgatattag      540
agcagctata cctaagcatt gttgggttaa gaatccatgg aagtctttga gttatgtcgt      600
cagagacgtc gctatcgtct ttgcattggc tgctggagct gcttacctca acaattggat      660
tgtttggect ctctattggc tcgctcaagg aaccatgttt tgggctctct ttgttcttgg      720
tcatgactgg taaacttaaa aaccctaact ttttcttctt tttctcctct gctttagtct      780
ccttttagcct ttgatttggt caactttgga tgattccaaa gaaccaatcg aacaaattgg      840
tctttatcca tatctcttca aatagcttta ggacataatt ggtctctcag gtaacaagct      900
gtcattatca tcatactcat catgttgcta gtagaccaac ccaattggca actgtttggt      960
ggttttgcaa ctgtgtaatc tgctttgaat tgtgaacaaa attattgatt tatgttgatt     1020
acattgcagt ggacatggta gtttctcaaa tgatccgaag ttgaacagtg tggtcgggtca     1080
tcttcttcat tcctcaatc tggtcccata ccatggctgg tgagttttgc ttccagacca     1140
ttcttctcta aaaccacttg cagaatctca tcttcttcat gtaaaaatat gactttgcag     1200
gagaattagt cacagaactc accaccagaa ccatggacat gttgagaatg acgaatcttg     1260

```

```

gcatcctgta agtcaaaaac gtattttttt ggttatcttg ttttagtcct gtgggtgtttc 1320
ttagatgcag ttttattaac tgtttctgta actgcagatg tctgagaaaa tctacaatac 1380
tttggaacag ccgactagat tcttttagatt tacactgcct ctctgtgatgc ttgcataccc 1440
tttctacttg gtaagaactc ctctatttgt tatggtaact taagctgccca caccaagtaa 1500
aaaagctcat gtctattctt ctgtttcagt gggctcgaag tccggggaaa aagggttctc 1560
attaccatcc agacagtgc ttgttctctc ctaaagagag aaaggatgtc ctcaattcta 1620
ctgcttggtg gactgcaatg gctgctctgc ttgtttgtct caacttcaca atcgggtccaa 1680
ttcaaatgct caaactttat ggaattcctt actgggtaat gcgccgctgt tactccccctg 1740
tttcagcctg agcaatttgt gtattatttc ctctgcctta ctcaaaaagg tttttatgtc 1800
aaatacagat aaatgtaatg tgggtggact ttgtgactta cctgcatcac catgggtcatg 1860
aagataagct tccttggtac cgtggcaagg taaaatacat attctctgct tccactgttc 1920
tttgactaca tcgctctttc ttttaagggt aaagccaact ggtgtgtaaa tctcatgatt 1980
ctcccaaac aggagtggag ttacctgaga ggaggactta caacattgga tcgtgactac 2040
ggattgatca ataacatcca tcatgatatt ggaactcatg tgatacatca tcttttcccg 2100
cagatcccac attatcatct agtagaagca gtaagtaaat tgaaagtaaa gactgtttgt 2160
gtttttggtg ttcatgctag tttccctgac tcttgctcca ctgttatgca gacagaagca 2220
gctaaaccag tattagggaa gtattacagg gagcctgata agtctggacc gttgccatta 2280
catttactgg aaattctagc gaaaagtata aaagaagatc attacgtgag cgacgaagga 2340
gaagttgtat actataaagc agatccaaat ctctatggag aggtcaaagt aagagcagat 2400
tgaaatgaag caggcttgag attgaagttt ttctatttc agaccagctg attttttgc 2460
tactgtatca atttattgtg tcaccacca gagagttagt atctctgaat acgatcgatc 2520
agatggaaac aacaaatttg tttgcgatac tgaagctata tataccata 2569

```

```

<210> 10
<211> 3879
<212> DNA
<213> Arabidopsis thaliana

```

```

<220>
<221> exon
<222> (780)...(1685)
<223> Exon 1

```

```

<221> exon
<222> (1761)...(2129)
<223> Exon 2

```

```

<221> exon
<222> (2207)...(2461)
<223> Exon 3

```

```

<221> exon
<222> (2544)...(2671)
<223> Exon 4

```

```

<221> exon
<222> (2762)...(2959)
<223> Exon 5

```

```

<221> exon
<222> (3088)...(3448)
<223> Exon 6

```

```

<400> 10
aaagatagta tttgttgata aatatgggga tatttatcct atattatctg tatttttctt

```

60

accatttttta	ctctattcct	ttatctacat	tacgtcatta	cactatcata	agatatttga	120
atgaacaaat	tcattgcacc	accagctata	ttaccctttt	ttattaaaaa	aaaacatctg	180
ataataataa	caaaaaaatt	agagaaatga	cgtcgaaaaa	aaaagtaaga	acgaagaaga	240
agtgttaaac	ccaaccaatt	ttgacttgaa	aaaaagcttc	aacgctcccc	ttttctcctt	300
ctccgctcgt	ctccgccgog	tcccaaatac	ccaattcctc	ctcttctccg	atcaattctt	360
cccaagtaag	cttcttcttc	ctcgattctc	tcctcagatt	gtttcgtgac	ttctttatat	420
atattcttca	cttcacagat	tttcttctgt	tgttgctgct	gatctcaaat	catagagatt	480
gattaaccta	attggtcttt	atctagtgtg	atgcacgtt	attaggaact	ttaaattaaag	540
atttaactgt	taatttcatg	attcggattc	gaattttact	gttctcgaga	ctgaaatatg	600
caacctattt	tttcgtaatc	gttggtgatc	aattcgaatt	ttcagaattt	atagcaattt	660
tgatgctcat	gatctgtcta	cgctacgttc	tcgtcgtaaa	tcgaagtgtg	taatgctatg	720
tgtttgttac	acaggtgtgt	gtatgtgtga	gagaggaaat	atagtgtaaa	aaattcataa	780
tggaagtctg	caattgtatt	gaaccgcaat	ggccagcgga	tgaattgtta	atgaaatacc	840
aatacatctc	cgatttcttc	attgcgattg	cgtatttttc	gattcctctt	gagttgattt	900
actttgtgaa	gaaatcagcc	gtgtttccgt	atagatgggt	acttgttcag	tttggtgctt	960
ttatcgttct	ttgtggagca	actcatctta	ttaacttatg	gactttcact	acgcattcga	1020
gaaccgtggc	gcttgtgatg	actaccgoga	agggtgttaac	cgctgttgct	tcgtgtgcta	1080
ctgctgtgat	gcttgttcat	attattcctg	atcttttgag	tgtaagact	cgggagcttt	1140
tcttgaaaaa	taaagctgct	gagctcgata	gagaaatggg	attgattcga	actcaggaag	1200
aaaccggaag	gcatgtgaga	atgttgactc	atgagattag	aagcacttta	gatagacata	1260
ctattttaaa	gactacactt	gttgagcttg	gtaggacatt	agctttggag	gagtggtcat	1320
tgtggatgcc	tactagaact	gggttagagc	tacagctttc	ttatacactt	cgcatcaac	1380
atcccgtgga	gtatacgggt	cctattcaat	taccggtgat	taaccaagtg	tttggtacta	1440
gtagggctgt	aaaaatatct	cctaattctc	ctgtggctag	gttgagacct	gtttctggga	1500
aataatgtct	aggggaggtg	gtcgtgtgga	gggttccgct	tctccacctt	tctaattttc	1560
agattaatga	ctggcctgag	ctttcaacaa	agagatatgc	tttgatgggt	ttgatgcttc	1620
cttcagatag	tgcaaggcaa	tggcatgtcc	atgagttgga	actcgttgaa	gtcgtcgctg	1680
atcaggtttt	acattgctga	gaatttctct	tctttgctat	gttcattgatc	ttgtctataa	1740
cttttctctc	cttattatag	gtggctgtag	ctctctcaca	tgctcgcgac	ctagaagagt	1800
cgatgcgagc	tagggacctt	ctcatggagc	agaatgttgc	tcttgatcta	gctagacgag	1860
aagcagaaac	agcaatccgt	gcccgcaatg	atttccctagc	ggttatgaac	catgaaatgc	1920
gaacaccgat	gcatgcgatt	attgcactct	cttccctact	ccaagaaacg	gaactaacc	1980
ctgaacaaag	actgatgggt	gaaacaatac	ttaaaagtag	taaccttttg	gcaactttga	2040
tgaatgatgt	cttagatctt	tcaaggtag	aagatggaag	tcttcaactt	gaacttggga	2100
cattcaatct	tcatacatta	tttagagagg	taacttttga	acagctctat	gtttcataag	2160
tttatactat	ttgtgtactt	gattgtcata	ttgaatcttg	ttgcagggtc	tcaatctgat	2220
aaagcctata	gcggttggtt	agaaattacc	catcacacta	aatcttgcac	cagatttgcc	2280
agaatttgtt	gttggggatg	agaaacggct	aatgcagata	atattaaata	tagttggtaa	2340
tgctgtgaaa	ttctccaaac	aaggtagtat	ctccgtaacc	gctcttgta	ccaagtcaga	2400
cacacgagct	gctgactttt	ttgtcgtgcc	aactgggagt	catttctact	tgagagtga	2460
ggttattatc	ttgtatcttg	ggatcttata	ccatagctga	aagtatttct	taggtcttaa	2520
ttttgatgat	tattcaaata	taggtaaaag	actctggagc	aggaataaat	cctcaagaca	2580
ttccaaagat	tttcaactaa	tttgctcaaa	cacaactctt	agcgacgaga	agctcgggtg	2640
gtagtgggct	tggcctcgcc	atctccaaga	ggtttgagcc	ttattaaaag	acgttttttt	2700
ccaacttttt	cttgtcttct	gtgttggtta	aagtttactc	ataagcggtt	aatatgacaa	2760
ggtttggtga	tctgatggag	ggtaacattt	ggattgagag	cgatgggtct	ggaaaaggat	2820
gcacggctat	ctttgatgtt	aaacttggga	tctcagaacg	ttcaaacgaa	tctaaacagt	2880
cgggcatacc	gaaagtcca	gccattcccc	gacattcaaa	tttcaactgga	cttaagggtc	2940
ttgtcatgga	tgagaacggg	ttagtataag	cttctcacct	ttctctttgc	aaaatctctc	3000
gccttacttc	ttgcaaatgc	agatattggc	gtttagaaaa	aacgcaaat	taatcttatg	3060
agaaaccgat	gattattttg	gttgagggtt	aagtagaatg	gtgacgaagg	gacttcttgt	3120
acaccttggg	tgcaagtgga	ccacgggtgag	ttcaaacgag	gagtggtctc	gagttgtgtc	3180
ccatgagcac	aaagtgggtc	tcattggacgt	gtgcatgccc	ggggtcgaaa	actaccaa	3240
cgctctccgt	attcacgaga	aattccaaaa	acaacgccac	caacggccac	tacttgtggc	3300
actcagtggg	aacactgaca	aatccacaaa	agagaaatgc	atgagctttg	gtctagacgg	3360

tgtgttgctc	aaacccgtat	cactagacaa	cataagagat	gttctgtctg	atcttctcga	3420
gccccgggta	ctgtacgagg	gcatgtaaag	gcatgggatg	cccatgccc	cagaggagta	3480
attccgctcc	cgccttcttc	tcccgtaaaa	catcggaagc	tgatgttctc	tggtttaatt	3540
gtgtacatat	cagagattgt	cggagcgttt	tgatgatata	cttaaaacag	aaaggggaata	3600
acaaaataga	aactctaaac	cggatgtgtg	ccgtggcgat	ttcggttata	gaggaacaag	3660
atgggtggtg	tataatcata	ccatttcaga	ttacatgttt	gactaatgtt	gtatccttat	3720
atatgtagtt	acattcttat	aagaatttgg	atcgagttaa	ggatgcttgt	tcggtgcatg	3780
tatgacattg	atgcagtatt	atggcgctcag	ctttgcgcgc	cttagtagaa	caacaacaat	3840
ggcgttactt	agtttctcaa	tcaacccgat	ctccaaaac			3879

<210> 11
 <211> 1200
 <212> DNA
 <213> Arabidopsis thaliana

<220>
 <221> CDS
 <222> (53) ... (1024)

cgttgctgtc	gaagttaggc	caagaaaccc	atttaaaaaa	aaagagagag	agatggagag	60
tttcccgatc	atcaatctcg	agaagcttaa	tggagaagag	agagcaatca	ctatggagaa	120
gatcaaagac	gcttgtgaaa	actggggctt	ctttgagtgt	gtgaaccatg	ggatttcact	180
cgagcttttg	gacaaagtgg	agaagatgac	caaggaacat	tacaagaagt	gcatggaaga	240
gagattcaag	gaatcgatta	agaacagagg	tcttgactct	cttcgctctg	aagtcaacga	300
cgttgactgg	gaatccactt	tctacctcaa	gcaccttccc	gtctctaata	tctccgatgt	360
ccctgatctc	gacgacgatt	acagaacggt	aatgaaagac	ttcgccggaa	agatagagaa	420
gttgctcgag	gagctactgg	atctgctgtg	cgagaatctc	gggttagaga	agggttatct	480
aaaaaagggt	ttttacgggt	cgaaaagacc	gacttttgga	accaaagtca	gcaattatcc	540
accttgtcct	aatccggacc	tagtcaaggg	tctccgagcc	cacaccgacg	ccggcggcat	600
catcctcctc	ttccaagacg	acaaaagtcag	tggacttcag	cttcttaaag	acggcgagtg	660
ggtcgatgtt	cctccgggta	agcattcaat	cgtcgttaat	ctcggcgatc	aacttgaggt	720
gataaccaat	gggaagtaca	agagtgtgga	acatagagtg	ctatctcaga	cagacggaga	780
aggaagaatg	tcgatcgcat	cattctataa	tccgggaagc	gactctgtta	tttttccggg	840
gccggagctg	atcggaaaaag	aagcagagaa	ggagaagaaa	gagaactatc	cgagatttgc	900
gtttgaagat	tacaggaaac	tctactctgc	tgtcaagttt	caggccaagg	aaccaagggt	960
tgaagccatg	aaagctatgg	agacaactgt	ggccaacaat	gttggaacct	tggccactgc	1020
gtgaatgata	tgttaactgg	taataaatat	atatatatat	atatatatag	tctttatata	1080
atgtcttaga	aacttgatta	ttcactatac	gaataatttt	gttcagtgtg	ttgtatgttt	1140
aagtgggtgaa	tgtgttatat	atgggaatta	atgttttctg	ttcgaaaaaa	aaaaaaaaaa	1200

<210> 12
 <211> 3438
 <212> DNA
 <213> Arabidopsis thaliana

<220>
 <221> exon
 <222> (1212) ... (1358)
 <223> Exon 1

<221> exon
 <222> (1461) ... (1592)
 <223> Exon 2

<221> exon
 <222> (1660) ... (1820)
 <223> Exon 3

<221> exon
 <222> (1909) ... (2893)
 <223> Exon 4

<400> 12

gttacttttc	aaatcttccc	tcataattata	tagccattga	tatcatagag	gatgtgagtt	60
ttaactttaat	atctaccgct	ttgaaactag	ctattttactt	aaatatgaat	tataatctag	120
tttaactacc	aaaaacatca	tatgggggaca	agaaaaagta	ataaaacgta	tggaaaaattt	180
tgtagatggt	ataaatggat	aattatttcaa	gtgataatct	atcactttga	tcttatctct	240
ttatccaatt	taattacttt	gtctctaagt	gatttgcttc	caaaatctaa	gtgtagtcta	300
tcctatt...	atcttatcct	atcatataat	cttctatata	tatgtgagtc	cgatgttgta	360
aagcgtacga	gagagagtaa	tgaagagtga	agtgttatat	tgttctctcg	tccacttcca	420
ctctctcttt	tatctcttac	ttacttcttc	gtaagatcat	tacatataat	aaataatatt	480
atztatqttt	gtgttatatt	taataacagt	aaaaagtttt	aaaacgttga	aaaaattagc	540
cgacatagaa	tacaaaagag	ggtagcatc	gggggagaaa	cgtggaccaa	catgatacac	600
cctccaaaat	agtcaccaag	ttgaaacatt	gacatgtttc	gctttttctt	ttctgtgtat	660
actttttttt	tctgtgggtc	acattattta	atatgtgtat	acaagcagct	attttacatg	720
gagatttcct	gtcggtatag	cgctctcatt	tctccatcgc	ttccactttt	ttcctatact	780
aatttgatct	aattaattca	tatgtcaaaa	cattaagaaa	atgaaactcg	taattcatac	840
ttgaatttaa	tagattaatt	aaaatgctat	ttattggcaa	aataaactcg	gtttatatct	900
aaatttttaga	atcactaaaa	ctttttgccc	aaaaaaaaat	aaaaataaat	cactaaaaaca	960
aaaaacaatc	aaaagaaaac	ccatgttggg	aaatcggata	atgaaaaata	ttagaatccc	1020
cgtcctttgt	gtattttggc	gtagcatgaa	actatataat	aaacatgcat	tcattcttag	1080
actctctgta	gcttatcaac	aacaacgcgc	tcgatctctc	tcagcctgtc	tgacaactct	1140
ttctctagtt	ctagagtttt	caattttattg	ttgagccttt	tattaaaaaa	aaaaaaacaa	1200
gaacaaaaga	aatgggtcaa	ttgtcaagaa	aagctacatg	caacagccat	ggccaagtct	1260
cttcgtattt	ccttggttgg	gaagagtacg	agaagaatcc	ttacgacgtt	accaagaacc	1320
ctcaaggcat	tatccagatg	ggtcttgccg	aaaatcaggt	aaacaaatat	tattcaacag	1380
catgtgatat	atatatactt	atgtatatca	tgacagagag	actaatttaa	agtatgttta	1440
atttttattgg	atttctgtag	ctatgctttg	atctactaga	gtcatggctt	gcacaaaaca	1500
cagacgcagc	ctgtttcaag	agagatggcc	agtctgtttt	ccgggaactc	gctctctttc	1560
aagactacca	tggcctctct	tccttcaaaa	atgtaagatt	attaattgta	tttatcaaat	1620
ttatttgtag	gttgctgata	ttgctcgaat	gattttcagg	cctttgctga	tttcatgtca	1680
gaaaatagag	gaaatcgagt	ttcttttgat	tcaacaaccc	ttgtgctcac	tgctggagcc	1740
acttcgcaa	acgagactct	aatgttttgt	cttgacagatc	cgggtgacgc	tttcttgctt	1800
cccacgccat	attatccagg	gttagtccac	tgtttgctta	cacgtaaaat	ttccatcatt	1860
cctacgaact	tgacttaact	aaaactcatg	tttatttttg	tacttcaggt	ttgataggga	1920
tctaaaaatgg	gaacccgctg	ttgagattgt	accaatccaa	agctcaagta	ctaaccgggt	1980
tcgcataacg	aaacttgac	tcgaagaagc	ctacgagcaa	gccaagaagc	ttgacctaaa	2040
cgtaaaagga	atactcatca	ccaacccatc	taaccctttg	ggtacgacaa	caacccaaac	2100
cgaaactcaac	attctatattg	atctcatcac	caagaataag	aatatacatt	tagtaagtga	2160
cgagatatat	tcgggcacag	tattcaactc	ttcagaattc	atcagcgtca	tggagattct	2220
aaaaaataat	caactcgaaa	acaccgatgt	tttgaaccga	gtccacattg	ttgttagctt	2280
atctaaagat	ctaggcctcc	ctgggttttag	agttggagcc	atctactcca	atgacaaaga	2340
tgctatctct	gccgctacaa	aaatgtcaag	tttcggcctt	gtctcctccc	agacacaata	2400
cctactatcc	tcattattat	ctgacaagaa	gttcactaag	aactacctta	gagagaacca	2460
aaaacggctc	aagaacagac	agagaaagct	cgtgttgggt	ctagaggcca	tcgggatcaa	2520
atgtctgaag	agtaatgcgg	gactcttttg	ttgggtcgac	atgagacctc	tccttagatc	2580
taaaacgttc	gaagcggaaa	tggatctttg	gaagaagatt	gtttacgaag	tgaagctcaa	2640
catctctcct	ggttcgtcgt	gccattgtga	agaaccgggt	tggtttagag	tttgtttcgc	2700
gaacatgatt	gatgagacat	taaagcttgc	tttaaagaga	ttgaagatgt	tggttgatga	2760

12

```

tgaaaaactca agtagaagat gccaaaagag taaaagcgaa agactaaacg gttcgaggaa 2820
gaagacgatg tcaaattgtct ctaactgggt tttccgacta tcgtttcacg accgtgaggc 2880
tgaggaacga tagtccggtt tttgttttga agttcttttt tttgttttcc cacacattgc 2940
aagtgtattct gtaatttttt ttatcacgag agagagtgtg aaaaaatgga aatgcaacgt 3000
gcttactctg atcctagatt ttagaaaacc gttgaagact tcttagagca agtccatcgg 3060
cagtttttaa tgggttttcta atgggttttct agctaattaa aagtccaaaa ttaaattgaaa 3120
acccaactaa ataattagga tccatcccaa tattagggtt tttggatggg ttttttagacg 3180
gcgacgtggg cgactgtgag tcgtcggaac acaaaaaaaaa tcacaacact catgttttcc 3240
tttttctctc cgtttttcac ttttttgttt tgtccgacgg ccggcgattc gaatcgattt 3300
gatctccggt gtatcgaaca tgaaatcggg agagaagagc caaatcatcg acgacttggt 3360
tcaccaattc cattcttcga accataactca tataagagtt tcttggttc tctctaaaac 3420
tcttctaatt ttctgata 3438

```

<210> 13

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Beneficial Oligonucleotide-Contains both DNA and
RNA

<400> 13

```

caggtcaagt gcaacgtagg atgattttta ucaaccuacg ttgcacuuga ccuggcgcggt 60
tttcgcgc 68

```

<210> 14

<211> 68

<212> DNA

<213> Artificial Sequence

<220>

<223> Beneficial Oligonucleotide-Contains Both DNA and
RNA

<400> 14

```

caggtcaagt gctacgtagg atgattttta ucaaccuacg tagcacuuga ccuggcgcggt 60
tttcgcgc 68

```

<210> 15

<211> 24

<212> DNA

<213> Jelly Fish

<400> 15

```

atggtgagca agggcgagga gctg 24

```

<210> 16

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Mutation

<400> 16
atggtgagca agggctagga gctg 24

<210> 17
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Mutation

<400> 17
atggtgagca agggcaggag ctgt 24

<210> 18
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Beneficial Oligonucleotide-Contains Both DNA and
RNA

<400> 18
gtgagcaagg gcgaggagct gttcattttu gaacagcucc tcgcccuugc ucacgcgcgt 60
tttcgcgc 68

<210> 19
<211> 68
<212> DNA
<213> Artificial Sequence

<220>
<223> Beneficial Oligonucleotide-Contains Both DNA and
RNA

<400> 19
tgagcaaggg ctggagctg ttcacttttg ugaacagcuc cgagcccuug cucagcgcgt 60
tttcgcgc 68

INTERNATIONAL SEARCH REPORT

Int. national application No.
PCT/US98/16267

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12N 15/82, 15/84, 15/82, 5/04; A01H 4/00

US CL : 536/23.6; 435/172.1; 800/278

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.6; 435/172.1; 800/278

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

BIOSIS, MEDLINE, AGRICOLA, CAPLUS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SPRINGER et al. Gene Trap Tagging of PROLIFERA, An Essential MCM2-3-5- Like Gene in Arabidopsis. Science. 12 May 1995, Vol. 268, pages 877-880. See the entire documentation.	1-45, 48-72
Y	SUNDARESAN et al. Patterns of Gene Action in Plant Development Revealed by Enhancer Trap and Gene Trap Transposable Elements. Genes Development. 1995, Vol. 9, No. 14, pages 1797-1810. See the entire documentation.	1-45, 48-72

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

A	document defining the general state of the art which is not considered to be of particular relevance	*T*	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
B	earlier document published on or after the international filing date	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O	document referring to an oral disclosure, use, exhibition or other means	*Z*	document member of the same patent family
P	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

13 OCTOBER 1998

Date of mailing of the international search report

30 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

OUSAMA M-FAIZ ZAGHMOUT

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16267

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 46-47
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1))(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/16267

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I. Claims 1-4, 8-30, 50-53 are drawn to a method of making localized mutation in a target gene.

Group II. Claims 5-7 are drawn to a method for making mutation using RNA segment contains at least 8 contiguous 2'-substituted Ribonucleotides.

Group III. Claims 31-45, 48-49 are drawn to a method of making localized, non-selectable mutation in a target gene.

Group IV. Claims 54-72 are drawn to a method of making specific mutation such as point mutation or frameshift mutation.

The inventions listed as groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The specific technical feature of group 1 is a method of making localized mutation in a target gene. Second product does not require the special technical features of group 1 because it entails to a method for making mutation using RNA segment contains at least 8 contiguous 2'-substituted Ribonucleotides, other than the ones claimed in group 1 and it does not require the particular DNA molecules of group 1. The third is a method of making localized, non-selectable mutation in a target gene, not required by group 1. The fourth is entails the making of point or frameshift mutation, does not require the special technical features of group I because it is drawn to specific rather than random mutation. The claims are not so linked by a special technical feature within the meaning of the PCT Rule 13.2 so as to form a single inventive concept, accordingly, the unity of invention is lacking among all groups.